

KDIGO - Controversies Conference



Oct 12-14, 2006 Amsterdam



Incidence and prevalence of CKD in Okinawa, Japan

Kunitoshi Iseki, MD


Dialysis Unit, University Hospital of The Ryukyus
Okinawa, Japan



Dialysis facility



Prevalence of ESRD in Japan



Screening Program in Japan

Law

- 1972 Employee (40 years and over)
- 1973 School children
- 1983 Urine test for elderly, annually
- 1997 10.67 millions people were tested

OGHMA (Okinawa General Health Maintenance Association)

Dipstick urinalysis (mandatory)

Serum creatinine (optional)


Standard analysis file

1983 N=106,182


1993 N=143,948

2003 N=154,019




Intermediate markers by GFR-stage




Metabolic syndrome



Overweight ($BMI \geq 25 \text{ kg/m}^2$)



GFR by Age-group



Japanese Society of Nephrology (JSN)

Membership	Number (ppm)
JSN Member	8747 (68.4)
Board Certified Nephrologist	2683 (21.0)
Attending Nephrologist	651 (5.1)

Activities

2006 CKD Initiative: JSN, JSdT, JSNP, JKF

2007 50th Annual Meeting of JSN

CKD Practice Guide Book

Asian Forum on CKD Initiative