

Full Review

Hypertension in dialysis patients: a consensus document by the European Renal and Cardiovascular Medicine (EURECA-m) working group of the European Renal Association–European Dialysis and Transplant Association (ERA-EDTA) and the Hypertension and the Kidney working group of the European Society of Hypertension (ESH)*

Pantelis A. Sarafidis¹, Alexandre Persu², Rajiv Agarwal³, Michel Burnier⁴, Peter de Leeuw⁵, Charles J. Ferro⁶, Jean-Michel Halimi⁷, Gunnar H. Heine⁸, Michel Jadoul⁹, Faical Jarraya¹⁰, Mehmet Kanbay¹¹, Francesca Mallamaci¹², Patrick B. Mark¹³, Alberto Ortiz¹⁴, Gianfranco Parati¹⁵, Roberto Pontremoli¹⁶, Patrick Rossignol¹⁷, Luis Ruilope¹⁸, Patricia Van der Niepen¹⁹, Raymond Vanholder²⁰, Marianne C. Verhaar²¹, Andrzej Wiecek²², Gregoire Wuerzner⁴, Gérard M. London²³ and Carmine Zoccali¹²

¹Department of Nephrology, Hippokration Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece, ²Pole of Cardiovascular Research, Institut de Recherche Expérimentale et Clinique, and Division of Cardiology, Cliniques Universitaires Saint-Luc, Université Catholique de Louvain, Brussels, Belgium, ³Department of Medicine, Indiana University School of Medicine and Richard L. Roudebush Veterans Administration Medical Center, Indianapolis, IN, USA, ⁴Service of Nephrology and Hypertension, Lausanne University Hospital, Lausanne, Switzerland, ⁵Department of Medicine, Maastricht University Medical Center, Maastricht and Zuyderland Medical Center, Geleen/Heerlen, The Netherlands, ⁶Department of Renal Medicine, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK, ⁷Service de Néphrologie-Immunologie Clinique, Hôpital Bretonneau, François-Rabelais University, Tours, France, ⁸Saarland University Medical Center, Internal Medicine IV–Nephrology and Hypertension, Homburg, Germany, ⁹Division of Nephrology, Cliniques Universitaires Saint-Luc, Université Catholique de Louvain, Brussels, Belgium, ¹⁰Department of Nephrology, Sfax University Hospital and Research Unit, Faculty of Medicine, Sfax University, Sfax, Tunisia, ¹¹Department of Medicine, Division of Nephrology, Koc University School of Medicine, Istanbul, Turkey, ¹²CNR-IFC, Clinical Epidemiology and Pathophysiology of Hypertension and Renal Diseases Unit, Ospedali Riuniti, Reggio Calabria, Italy, ¹³Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK, ¹⁴IIS-Fundacion Jimenez Diaz, School of Medicine, University Autonoma of Madrid, FRIAT and REDINREN, Madrid, Spain, ¹⁵Department of Cardiovascular, Neural, and Metabolic Sciences, San Luca Hospital, Istituto Auxologico Italiano and Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy, ¹⁶Università degli Studi and IRCCS Azienda Ospedaliera Universitaria San Martino-IST, Genova, Italy, ¹⁷INSERM, Centre d'Investigations Cliniques Plurithématique 1433, UMR 1116, Université de Lorraine, CHRU de Nancy, F-CRIN INI-CRCT Cardiovascular and Renal Clinical Trialists, and Association Lorraine de Traitement de l'Insuffisance Rénale, Nancy, France, ¹⁸Hypertension Unit & Institute of Research i + 12, Hospital Universitario 12 de Octubre, Madrid, Spain, ¹⁹Department of Nephrology and Hypertension, Universitair Ziekenhuis Brussel - VUB, Brussels, Belgium, ²⁰Nephrology Section, Department of Internal Medicine, Ghent University Hospital, Gent, Belgium, ²¹Department of Nephrology and Hypertension, University Medical Center Utrecht, The Netherlands, ²²Department of Nephrology, Transplantation and Internal Medicine, Medical University of Silesia in Katowice, Katowice, Poland and ²³Manhes Hospital and FCRIN INI-CRCTC, Manhes, France

ABSTRACT

In patients with end-stage renal disease (ESRD) treated with haemodialysis or peritoneal dialysis, hypertension is common and often poorly controlled. Blood pressure (BP) recordings obtained before or after haemodialysis display a J- or U-shaped association with cardiovascular events and survival, but this most likely reflects the low accuracy of these measurements and the peculiar haemodynamic setting related to dialysis treatment. Elevated BP detected by home or ambulatory BP monitoring is clearly associated with shorter survival. Sodium and volume excess is the prominent mechanism of hypertension in dialysis patients, but other pathways, such as arterial stiffness, activation of the renin-angiotensin-aldosterone and sympathetic nervous systems, endothelial dysfunction, sleep apnoea and the use of erythropoietin-stimulating agents may also be involved. Non-pharmacologic interventions targeting sodium and volume excess are fundamental for hypertension control in this population. If BP remains elevated after appropriate treatment of sodium and volume excess, the use of antihypertensive agents is necessary. Drug treatment in the dialysis population should take into consideration the patient's comorbidities and specific characteristics of each agent, such as dialysability. This document is an overview of the diagnosis, epidemiology, pathogenesis and treatment of hypertension in patients on dialysis, aiming to offer the renal physician practical recommendations based on current knowledge and expert opinion and to highlight areas for future research.

Keywords: blood pressure, end-stage renal disease, haemodialysis, hypertension, peritoneal dialysis

INTRODUCTION

In patients with end-stage renal disease (ESRD) receiving renal replacement therapy with haemodialysis or peritoneal dialysis, hypertension is very common and often inadequately controlled [1]. Elevated blood pressure (BP), particularly when recorded outside of the dialysis unit with home or ambulatory BP monitoring, is directly associated with shorter survival [2–4]. Sodium and volume excess appear to be the most important causes of hypertension in dialysis patients; therefore, non-pharmacologic strategies such as dietary sodium restriction, individualized dialysate sodium prescription and gradual dry-weight reduction should be the initial therapeutic approaches to control BP [5, 6], but this approach is often not adequately implemented [7, 8]. In patients who remain hypertensive after management of sodium and volume excess, pharmacological therapy is recommended

to achieve BP control, taking into account the pharmacologic characteristics of each antihypertensive drug [5, 6, 9].

This is a document prepared by experts from the European Renal and Cardiovascular Medicine (EURECA-m) working group of the European Renal Association–European Dialysis and Transplant Association (ERA-EDTA) and the Hypertension in the Kidney working group of the European Society of Hypertension (ESH). It aims to summarize current knowledge on the diagnosis, epidemiology, pathogenesis and treatment of hypertension in ESRD patients on dialysis. As far as treatment is concerned, we discuss both non-pharmacological and pharmacological strategies to manage hypertension. This document mainly presents evidence from patients receiving maintenance haemodialysis treatment, because most of the current knowledge is derived from studies of this category of patients. Data from the few relevant studies concerning peritoneal dialysis patients are also discussed.

DIAGNOSIS OF HYPERTENSION IN DIALYSIS PATIENTS

According to the 2004 National Kidney Foundation Kidney Disease Outcomes Quality Initiative guidelines [10], hypertension in haemodialysis patients is diagnosed when pre-dialysis BP is >140/90 mmHg or when post-dialysis BP is >130/80 mmHg [10]. However, the diagnosis of hypertension using conventional peridialytic BP recordings may be problematic for several reasons [11, 12]. Pre- and post-dialysis BPs are recorded by dialysis unit staff, often without the necessary attention to the standardization of the technique of BP measurement and the prerequisites for objective office BP recordings [13]. BP measurements pre-, during and post-dialysis are not made for diagnostic reasons but to exploit a major haemodynamic metric like BP in order to assess cardiovascular stability before, during and immediately after the dialysis procedure. Thus, using these readings to diagnose hypertension, assess the success of antihypertensive treatment or examine future cardiovascular risk is inherently problematic. Several factors may lead to inaccurate BP pre- and post-dialysis readings, such as the white coat effect, limited time for relaxation (patient impatience to start dialysis and leave the unit quickly), fear or anxiety for correct arteriovenous fistula needling, previous bilateral upper limb attempts of arteriovenous fistulae and the unknown validity of most oscillometric devices attached to commercially available haemodialysis machines. Furthermore, truly high BP variability (pre- to post-dialysis and day by day variability) in response to fluctuations in volume status and other parameters during the

intra- and interdialytic periods is another important issue that complicates the accurate diagnosis of hypertension [14]. The typical pattern of haemodynamic response to ultrafiltration is BP decrease from pre- to post-dialysis; the magnitude of intradialytic BP reduction is, for the most part, related to the magnitude and rate of volume withdrawal during dialysis. The converse phenomenon is observed during the interdialytic interval [15], and several studies show that interdialytic weight gain is closely associated with higher pre-dialysis BP [16]. The poor diagnostic accuracy of peridialytic BP recordings is supported by a meta-analysis showing that both pre- and post-dialytic BP readings provide imprecise estimates of the mean interdialytic BP recorded with 44-h ambulatory BP monitoring (ABPM) [17]. Furthermore, in comparison with interdialytic BP recordings, peridialytic BP recordings have a weaker prognostic relationship with mortality in haemodialysis patients [2, 3, 11]. It must be noted that it is not known whether peridialytic measurements following a standardized technique would exhibit stronger prognostic associations with outcome; preliminary evidence suggests that this is not very likely since, even when peridialytic BP is recorded with a standardized protocol, it relates poorly to 44-h ABPM values [18].

Due to the reasons described above, the rate of errors in the diagnosis of hypertension when using peridialytic BP measurements is unacceptably high [19]. The proportions of chronic kidney disease (CKD) patients with white-coat and masked hypertension are reported to be around 30 and 7%, respectively, but are suggested to be much higher in people receiving dialysis [14, 20–22]. An alternative can be the use of an average of intradialytic BP measurements; in one study, a median intradialytic cut-off BP of 140/90 mmHg during a mid-week dialysis session provided greater sensitivity and specificity in detecting interdialytic hypertension compared with pre- and post-dialysis BP measurements [23]. Yet, BP measurements obtained outside of dialysis units are still needed to reliably diagnose hypertension among dialysis patients. Home BP monitoring is widely applied and is strongly recommended by international guidelines for the diagnosis and management of hypertension in the general population [24]. Compared with BP recordings obtained pre- or post-dialysis, home BP exhibits stronger associations with mean 44-h ambulatory BP [18, 20]. In the Dry-Weight Reduction in Haemodialysis Patients (DRIP) trial, changes in home BP after 4 and 8 weeks of dry-weight probing (i.e. supervised gradual dry-weight reduction) were closely associated with changes in 44-h ambulatory BP; in contrast, pre- and post-dialysis BP recordings failed to detect the changes in ambulatory BP caused in response to dry-weight reduction [25]. Moreover, home BP was shown to have high short-term reproducibility from one week to the next [20], in contrast to the high variability and poor reproducibility of conventional peridialytic BP recordings [14]. Furthermore, home BP exhibits stronger associations with indices of target organ damage [26–28] and represents a more powerful predictor of future cardiovascular events or all-cause mortality compared with the BP measurements obtained within dialysis units [2, 3, 11]. It is important to note that interdialytic BP recordings maintain their strong prognostic association with cardiovascular outcomes, even when a small number (i.e. six) randomly selected measurements are used to

assess the interdialytic BP burden [29]; thus, the location and time-frame covered (and not the number of BP recordings) is the major factor determining the strong prognostic significance of interdialytic ambulatory BP measurements, although the timing of BP recordings may be relevant for reproducibility [30]. The notion that home BP is useful to guide the management of hypertension in dialysis patients is supported by a pilot study that randomized 65 hypertensive haemodialysis patients to have their antihypertensive drug therapy adjusted either on the basis of routine pre-dialytic BP or with home BP monitoring. Over a mean follow-up of 6 months, a significant reduction in interdialytic ambulatory BP of 9/7 mmHg was documented in the home BP-guided group, but not in the pre-dialytic BP-guided group [31]. Similar results were registered in another small randomized trial in haemodialysis patients [32]. However, one important aspect is for future studies to gather data in order to provide patients with a precise protocol on when and how often home BP measurements should be performed, as has been done for hypertensive patients in the general population [31].

Many authors suggest that ABPM is the ‘gold standard’ method for diagnosing hypertension in patients receiving dialysis [5, 11, 33, 34]. The superiority of this approach over conventional peridialytic BP measurements is strongly supported by comparative studies showing that mean 44-h interdialytic BP is more strongly associated with the presence of target organ damage [such as echocardiographic left ventricular hypertrophy (LVH)] [26]. In addition, observational studies clearly suggest that ABPM predicts all-cause and cardiovascular mortality better than peridialytic BP [2, 4, 11]. The use of ABPM also has the advantage of recording BP during the night, providing additional information on the circadian variation of BP; the presence of a non-dipping nocturnal BP pattern is very common among dialysis patients and has been associated with LVH [35] and increased risk of all-cause and cardiovascular mortality [36]. The high prevalence of non-dipping and nocturnal hypertension among dialysis patients [12] suggests that the application of ABPM for the diagnosis and treatment of hypertension is more compelling than in the general population, where ABPM has already been strongly recommended by an *ad hoc* ESH working group [37], NICE guidelines [38] and the US preventive service [39]. The thresholds to define hypertension using home and ambulatory BP monitoring [11] are summarized in Box 1. Of note, when neither ABPM nor home BP measurements are applicable in dialysis patients, the diagnosis and management of hypertension can be made on the basis of office BP measurements taken during the interdialytic interval, as a recent study suggested that, in contrast to pre-dialysis BP, which has a U-shaped relationship with mortality, in the same patients the average of three office measurements (obtained by trained personnel from patients in the sitting position after at least 5 min of quiet rest) is almost linearly related to this risk [42]. The threshold of office BP (140/90 mmHg) recommended by current guidelines for the definition of hypertension in CKD patients [43] can also be extended to haemodialysis patients; however, it has to be noted that the issue of the optimal BP in CKD is controversial [43] and could be re-examined in the near future in view of recent evidence [44].

Box 1: Diagnosis of hypertension in dialysis patients

Hypertension in dialysis patients should be defined on the basis of home BP or ABPM measurements. Thresholds and methods proposed by the ASH/ASN [5], the EURECA-m working group of ERA-EDTA [11] and the relevant ESH Guidelines [24, 40, 41] can be used as follows:

- Home BP in haemodialysis: an average BP $\geq 135/85$ mmHg for measurements collected in the morning and in the evening over 6 non-dialysis days (covering a period of 2 weeks). Measures should be performed in a quiet room, with the patient in seated position, back and arm supported, after 5 min of rest and with two measurements per occasion taken 1–2 min apart.
- Home BP in peritoneal dialysis: an average BP $\geq 135/85$ mmHg over 7 consecutive days with measurements collected as above.
- ABPM in haemodialysis: an average BP $\geq 130/80$ mmHg over 24-h monitoring during a mid-week day free of haemodialysis. Whenever feasible, ABPM should be extended to 44 h, that is, covering a whole mid-week dialysis interval.
- ABPM in peritoneal dialysis: an average BP $\geq 130/80$ mmHg over 24-h monitoring.
- For haemodialysis patients, no recommendation can be made on the basis of pre- or post-dialysis BP. When neither ABPM nor home BP measurements are available in these patients, the diagnosis can be made on the basis of office BP measurements taken in a mid-week day free of haemodialysis, that is, the average of three measurements with 1–2 min intervals obtained in the sitting position by trained personnel after at least 5 min of quiet rest. The threshold of office BP $\geq 140/90$ mmHg recommended by current guidelines for the definition of hypertension in CKD patients can be used for haemodialysis patients.
- For peritoneal dialysis patients, office BP $\geq 140/90$ mmHg obtained as described immediately above can be used for the diagnosis of hypertension.

BP, blood pressure; ABPM, ambulatory blood pressure monitoring; ASH, American Society of Hypertension; ASN, American Society of Nephrology; EURECA-m, European Renal and Cardiovascular Medicine working group; ERA-EDTA, European Renal Association–European Dialysis and Transplant Association; ESH, European Society of Hypertension.

Despite the above advantages, ABPM is still perceived as a technique with limited applicability in dialysis patients. This reservation is partly due to the fact that a substantial number of the studies using ABPM in dialysis patients conducted to date were performed in a single American academic haemodialysis unit [2, 15, 26], but also to the fact that ABPM is believed to be uncomfortable and inconvenient in a group of patients with a high treatment burden, including a high proportion of sleep disturbances, especially when applied for 48 h. Furthermore, accurate ABPM readings could be challenging in patients with bilateral upper limb attempts of arteriovenous fistulae for dialysis access [11, 45]. The fact that ABPM is not reimbursed in many countries is another obstacle to its wider use in haemodialysis. However, additional research is needed to define the acceptability of ABPM from the patients', the best BP thresholds to define hypertension, which may be different from those of the general population because of the continuous shifts of volume and other factors, the optimum frequency of its use and the cost-effectiveness of ABPM in the dialysis population. Until

ongoing studies investigating these issues become available, home BP monitoring appears to be a simpler and more efficient approach to measure BP and make therapeutic decisions in dialysis patients [19].

In contrast to the typical decline in BP during dialysis, in ~ 10 – 15% of dialysis patients, BP exhibits a 'paradoxical' intradialytic elevation [46, 47]. Although this abnormal pattern of intradialytic haemodynamic response has been long recognized, the exact definition of intradialysis hypertension is still a matter of debate. For example, in some studies, intradialysis hypertension is defined as a rise of at least 15 mmHg in mean BP during dialysis [48], while others define it as a rise of at least 10 mmHg in systolic BP (SBP) during dialysis or immediately post-dialysis in a certain number (most commonly the last three or four out of the last six) of dialysis treatments [46, 47, 49, 50] or with the use of the regression of all intradialytic BP measurements over time with a slope greater than zero [51]. A case-control study that compared the interdialytic BP profile of 25 patients with intradialysis hypertension with that of 25 age- and sex-matched controls with normal intradialytic haemodynamic response [52] made the important observation that intradialysis hypertension is a phenomenon superimposed upon background interdialytic hypertension, as patients with intradialysis hypertension had higher 44-h interdialytic BP than controls. Of note, patients with intradialysis hypertension also had a gradual BP decline during the first 24 h after dialysis, which contrasted with the (typical) gradual increase from post-dialysis onwards in patients without intradialytic hypertension.

PREVALENCE OF HYPERTENSION IN THE HAEMODIALYSIS POPULATION BY THE VARIOUS METRICS AND DEFINITIONS

The estimates of the prevalence, treatment and control of hypertension among patients on chronic dialysis are highly variable. This variability arises, in large part, from differences in the definitions used to diagnose hypertension and in the setting of BP measurement (i.e. routine peridialytic BP recordings or interdialytic ABPM) in various studies [1, 53–56].

Office or peridialytic BP recordings

Hypertension is highly prevalent among patients with CKD who are not yet on dialysis. In a cross-sectional analysis of 10 813 CKD patients participating in the Kidney Early Evaluation Programme in the USA, hypertension (defined as BP $\geq 130/80$ mmHg or use of antihypertensive drugs) was detected in 86.2% of the overall study cohort; prevalence of hypertension exhibited a stepwise increase with advancing stage of CKD, increasing to 95.5% (or 91% with the use of a 140/90 threshold) in participants with stage 4 and 5 CKD [57]. A study of patients with pre-dialysis CKD that were followed in a low-clearance clinic [mean estimated glomerular filtration rate (eGFR) 14.5 mL/min/1.73 m²] similarly showed the prevalence of hypertension to be 95% [58], indicating that almost all CKD patients are hypertensive just before the initiation of renal replacement therapy.

Initiation of dialysis may have a substantial impact on the management of hypertension, as dialysis represents a potent

Table 1. Prevalence, treatment and control of hypertension in haemodialysis patients

Author	Year	N	Definition of hypertension	Prevalence of hypertension (%)	BP treatment among hypertensives (%)	BP control among hypertensives (%)
Salem [55]	1995	649	Pre-haemodialysis MAP ≥ 114 mmHg or use of antihypertensive agents	71.9	81.5	48.6
Rahman <i>et al.</i> [60]	1999	489	Pre-haemodialysis SBP ≥ 140 mmHg and/or DBP ≥ 90 mm	87.7	93.2	71.1
Agarwal <i>et al.</i> [1]	2003	2535	1-week average pre-haemodialysis SBP > 150 mmHg and/or DBP > 85 mmHg, or use of antihypertensive agents	85.8	88.4	30.3
Agarwal [56]	2011	369	44-h interdialytic ambulatory SBP ≥ 135 mmHg and/or DBP ≥ 85 mmHg or use of antihypertensive medications	82	89	38

BP, MAP, blood pressure; MAP, mean arterial pressure; SBP, diastolic blood pressure; DBP, systolic blood pressure.

therapeutic tool to remove sodium and fluid excess and improve BP control. Thus, hypertension prevalence in dialysis patients may appear lower than in those with pre-dialysis CKD. However, hypertension prevalence after initiation of dialysis depends on the clinical policies adopted in each dialysis unit. In some units, where long dialysis and strict control of salt intake are prescribed, hypertension has a lower prevalence than in those where such clinical policies are not applied [59]. However, increasing dialysis time to >4 h may be not feasible due to a number of factors, including limited facility and staff resources, patient preferences and others.

In epidemiology studies in haemodialysis patients in the USA that used different ways to define hypertension, the prevalence of hypertension ranged between 72 and 88% of the total population studied (Table 1). Despite the high proportion of hypertensive patients using antihypertensive medications, the amount of those that had their BP under control was low in the majority of these studies, that is, roughly between 30 and 50% [1, 55, 60]. Information on hypertension prevalence in dialysis patients in countries other than the USA is limited. In studies made within the framework of the Dialysis Outcomes and Practice Patterns Study (DOPPS), the prevalence of hypertension was very high and rose over time in all countries. In the last of these surveys (2011), hypertension prevalence ranged from 78% in Japan to 96% in Germany [61].

Interdialytic ambulatory BP monitoring

When estimated by the 'gold standard' method of 44-h interdialytic ABPM, with hypertension defined as average SBP ≥ 135 mmHg and/or diastolic BP (DBP) ≥ 85 mmHg or the use of antihypertensive medications, the prevalence of hypertension was 82% in a population of 369 predominantly African American patients who received haemodialysis treatment in units affiliated with an American university [56]. Eighty-nine percent of hypertensives were treated with antihypertensive drugs, but the rate of 44-h BP control (i.e. patients with average BP below the above thresholds) was as low as 38% [56]. Poor hypertension control in this study was associated with higher number of antihypertensive drugs and fluid overload as measured by the inferior vena cava diameter in expiration [62].

Apart from this study in African Americans, no large surveys reporting hypertension prevalence in dialysis patients based on ABPM have been performed in other ethnicities and countries to date.

BP AND THE RISK FOR CARDIOVASCULAR EVENTS AND DEATH IN HAEMODIALYSIS PATIENTS

The relationship of BP with all-cause and cause-specific mortality in haemodialysis patients is a controversial issue. Several studies have shown that in the BP range (i.e. SBP 110–180 mmHg) in which the event risk increases substantially with BP increase in the general population, there is either no relationship or a U-shaped association of pre- or post-dialysis SBP and DBP with all-cause and cardiovascular mortality [63–66], a phenomenon described as 'reverse epidemiology of hypertension' in the dialysis population. Some studies suggested that low BP in haemodialysis is associated with early mortality and deaths of primarily non-cardiac origin, indicating poor physiological reserve and frailty due to comorbid conditions (e.g. terminal cancer or congestive heart failure) to be the underlying factors of mortality [67]. However, this flat or U-shaped association raised substantial concerns about whether BP-lowering, as a whole, is a strategy associated with benefits for these patients [68]. More recent observations support that this phenomenon is due to the inadequacy of peridialytic BP recordings *per se* to describe the true BP load, rather than a true flat or U-shaped relationship of BP with cardiovascular morbidity and mortality. Of note, a study of more than 44 000 haemodialysis patients in the USA suggested post-dialysis pulse pressure (PP) to be associated with higher risk of mortality (12% higher risk for every 10 mmHg increase in PP), whereas post-dialysis SBP displayed an inverse relationship with risk [69]. In another cohort of 11 142 haemodialysis patients, high post-dialysis SBP and low pre- and post-dialysis DBP were associated with mortality, again implicating high PP as a causal factor [70]. Further to that, a recent analysis of 24 525 patients from the DOPPS study indicated that the U-shape between BP and mortality was

mostly observed for SBP (pre-dialysis SBP <130 mmHg or ≥ 160 mmHg was associated with higher mortality), but not for DBP, where a higher mortality rate was only observed in patients with pre-dialysis DBP <60 mmHg, suggesting that increased PP/arterial stiffness and/or comorbid conditions may be responsible for these associations [71].

In contrast to the unclear association of peridialytic BP recordings with all-cause and cardiovascular mortality, prospective cohort studies have shown that interdialytic BP recorded either at home or by ABPM associates more clearly with mortality and cardiovascular events, as it is also documented for the general population. In a group of 57 treated hypertensive haemodialysis patients prospectively followed for a mean period of 34.4 ± 20.4 months, Amar *et al.* [4] showed elevated 24-h ambulatory PP [relative risk (RR): 1.85 for each 10 mmHg increase in PP; 95% confidence interval (95% CI): 1.28–2.65], as well as elevated nocturnal SBP (RR: 1.41 for each 10 mmHg increase in nocturnal SBP; 95% CI: 1.08–1.84), to be independently associated with increased risk of cardiovascular mortality. In a larger study by Tripepi *et al.* [36], in 168 non-diabetic haemodialysis patients, nocturnal BP burden (as estimated by the night/day ratio) was a direct predictor of a surrogate end point such as LVH, as well as of cardiovascular events and death. A clear association between average interdialysis BP, as measured by home BP or ABPM, and mortality was described by Alborzi *et al.* in a cohort of 150 haemodialysis patients, while no such relationship was evident for pre-dialysis BP measurements (Figure 1) [3]. In the largest study performed so far, undertaken in 326 mainly African American patients, patients in the higher quartiles of home and 44-h ambulatory SBP exhibited an excessive risk of mortality, which was independent of other risk factors over 32 months of follow-up [2].

Additional support for the notion that interdialytic BP recordings have closer association with outcomes is provided by a recent prospective analysis of patients participating in the Chronic Renal Insufficiency Cohort study [42]. The prognostic association of SBP with all-cause mortality was assessed at three different time-points in this prospective cohort: (i) when

participants had stage 4 CKD (eGFR <30 mL/min/1.73 m²); (ii) when participants-initiated haemodialysis and dialysis-unit BP measurements were available; and (iii) when incident haemodialysis patients had an out-of-dialysis BP measurement obtained during a pre-specified follow-up visit at home. SBP had no association with mortality among participants not yet on dialysis. In accordance with earlier reports from other cohorts of haemodialysis patients, dialysis-unit SBP provided a U-shaped association with mortality. In contrast, a direct linear association between SBP and all-cause mortality was evident when BP measurements were obtained outside the unit [hazard ratio (HR): 1.26 for each 10 mmHg higher SBP; 95% CI: 1.14–1.40] [42].

EPIDEMIOLOGY OF HYPERTENSION IN PATIENTS TREATED WITH PERITONEAL DIALYSIS

The prevalence of hypertension among patients on peritoneal dialysis was evaluated in a cross-sectional study conducted on 504 patients in 27 peritoneal dialysis centres of the Italian Co-operative Peritoneal Dialysis Study Group [72]. Valid ambulatory BP measurements were obtained in 414 patients (82%). Using the WHO/ISH 1999 definition of hypertension (SBP >140 or DBP >90 mmHg, or use of antihypertensive treatment), the prevalence of hypertension was 88%. When hypertension was defined using a BP load of >30% of values >140/90 at day-time or >120/80 at night-time during 24-h ABPM, the estimated prevalence of hypertension was lower (69%); however, the actual ability of BP load to identify a hypertensive condition has been questioned [37]. The average 24-h BP in this study was $139 \pm 19/81 \pm 11$ mmHg, again suggesting that, if the currently proposed definition of average SBP ≥ 135 and/or DBP ≥ 85 mmHg in ABPM or antihypertensive treatment [5] was used instead, hypertension prevalence would also exceed 70–80% [72]. Of note, 53% of patients in this study were non-dippers and an additional 9% were reverse-dippers. Small studies comparing the ambulatory BP profile between patients treated with automated peritoneal dialysis versus continuous ambulatory peritoneal dialysis showed that the average 24-h BP did not differ between the treatment modalities [73, 74]. Other studies have described an association between BP and peritoneal transport status; patients with high peritoneal transport (reflecting poor peritoneal ultrafiltration) have higher BP levels during both day-time and night-time periods, as well as higher left ventricular (LV) mass index compared with 'low transporters', and this difference most likely reflects volume overload triggered by high peritoneal transport and subsequent decreased ultrafiltration capacity in the first group [75]. Volume overload is frequently more marked in peritoneal dialysis than in haemodialysis patients [76] and these patients require antihypertensive drugs more frequently (65%) than haemodialysis patients (38%; $P < 0.001$). The detrimental role of volume excess in patients maintained for too long on peritoneal dialysis is well described [77]. In this regard, a strict volume control policy could reduce the need for antihypertensive medication in peritoneal dialysis patients.




FIGURE 1: Hazard ratios for all-cause mortality for quartiles of pre-dialysis, post-dialysis, home and ambulatory systolic blood pressure (BP). Higher levels of home BP and ambulatory BP were significantly associated with mortality, whereas pre- and post-dialysis BP was not. P-values are those reported for linear trend. HD, haemodialysis; Q, quartile. Reproduced with permission from Alborzi *et al.* [3]

Given the more continuous nature of renal replacement therapy and the absence of cyclic variations in volume status and several other metabolic parameters in patients receiving peritoneal dialysis, it has been hypothesized that BP control and BP diurnal variation may differ substantially between patients treated with peritoneal dialysis or haemodialysis. However, only two studies have tested this to date. One [75] compared the 44-h BP profile of 22 haemodialysis patients with that of 24 patients treated with continuous ambulatory peritoneal dialysis. Mean 44-h SBP and DBP was no different between the two dialytic modalities; however, in haemodialysis patients, night-time BP recorded on the dialysis-off day was significantly higher and day-time BP recorded on the dialysis-on day was significantly lower than in continuous ambulatory peritoneal dialysis patients [75]. Another study, including 33 haemodialysis and 27 peritoneal dialysis patients, showed that diurnal BP pattern (i.e. dipping status) did not differ between the two modalities over a ~48-h BP recording period, but that average ambulatory SBP (142.1 ± 16.3 versus 130.4 ± 17.1 mmHg; $P < 0.01$) and SBP loads ($54 \pm 29\%$ versus $30 \pm 31\%$; $P < 0.01$) were higher in those receiving haemodialysis [78]. Overall, the above studies are small and largely inconclusive; methodologically rigorous comparisons between haemodialysis and peritoneal dialysis patients are missing and seem rather unfeasible.

PATHOPHYSIOLOGY OF HYPERTENSION IN DIALYSIS PATIENTS

Increase in cardiac output, peripheral vascular resistance or both may result in sustained BP elevation among patients on dialysis. Undoubtedly, sodium and volume overload are considered the prominent pathogenic mechanisms. A number of non-volume mediated pathways, such as activation of the renin–angiotensin–aldosterone and sympathetic nervous systems, structural arterial wall alterations related to the long-term arteriosclerotic process, endothelial dysfunction, inflammation, sleep apnoea and use of particular medications like erythropoietin-stimulating agents, are also reported to play an important role in the complex pathogenesis of hypertension in these individuals (Box 2) [79].

Volume overload

In patients with ESRD, even when residual renal function is preserved, the sodium and fluid excretory capacity is

Box 2: Main pathogenic mechanisms of hypertension in dialysis patients

- Sodium and volume overload.
- Increased arterial stiffness.
- Activation of the sympathetic nervous system.
- Activation of the renin–angiotensin–aldosterone system.
- Endothelial dysfunction (i.e. imbalance between endothelium-derived vasodilators and vasoconstrictors).
- High prevalence of sleep apnoea.
- Use of recombinant erythropoietins.

substantially impaired. Thus, sodium retention and volume overload is very common and often not easily identifiable. Moreover, ESRD patients have the highest sodium sensitivity of BP [80, 81]. It is now well documented that, in addition to classical osmotic volume expansion, sodium retention may occur in the form of osmotically inactive sodium in the connective tissue and skin, where sodium accumulates linked to glycosaminoglycans [82]. Non-osmotic sodium retention triggers local macrophage recruitment; macrophages sense the hypertonic electrolyte accumulation in the skin and activate the tonicity-responsive enhancer-binding protein to initiate the secretion of vascular endothelial growth factor (VEGF), which enhances electrolyte clearance via cutaneous lymph vessels and increases endothelial nitric oxide (NO) synthase expression in blood vessels. Deletion of tonicity-responsive enhancer-binding protein in monocytes or blockade of lymph–endothelial VEGF receptor inhibits lymphogenesis, promotes endothelial dysfunction and increases BP in mice in response to salt loading [83]; that is, hypertension is promoted by mechanisms different to those traditionally ascribed to iso-osmotic retention. In haemodialysis patients, sodium and water in skin and muscle are increased and VEGF is reduced when compared with age-matched healthy individuals; these phenomena may also contribute to hypertension [84]. Due to sodium and fluid accumulation, BP steadily increases in proportion to weight gain during the interdialytic interval, a phenomenon superimposed on BP circadian variation [85]. The interdialysis increase in BP is not limited to brachial BP but extends to other critical haemodynamic parameters like aortic BP [86], and the peripheral and central BP burden is accentuated during the long dialysis interval (Figure 2), again in proportion to fluid overload [87, 88]. Until fluid and sodium overload is removed during dialysis, a rise in peripheral vascular resistance will sustain hypertension in such individuals.

Arterial stiffness increase

Patients with ESRD display a premature increase in arterial stiffness due to a combination of factors, mainly as a result of




FIGURE 2: Changes in aortic blood pressures, wave reflections and arterial stiffness parameters between the first and the second interdialytic day, Δ [day (2) – day (1)], in comparison with relevant changes between the second and the third interdialytic day, Δ [day (3) – day (2)]. Reprinted with permission from Koutroumbas *et al.* [87]. AoSBP, aortic systolic blood pressure; AoDBP, aortic diastolic blood pressure; AoPP, aortic pulse pressure; MBP, mean blood pressure; AIx, augmentation index; PWV, pulse wave velocity.

disturbed calcium phosphate homeostasis [89]. In dialysis, arterial stiffness, assessed by aortic pulse wave velocity (PWV), determines the patterns and rhythms of BP recorded over the interdialytic period [89–91]. Agarwal and Light [90] analysed 11 833 interdialytic BP measurements from 125 haemodialysis patients and showed that log of PWV was related to BP in a linear relationship (each log increase in PWV was associated with 20.3, 7.2 and 12.8 mmHg increases in SBP, DBP and PP, respectively). Increasing PWV also blunted the circadian amplitude of SBP and PP. In a *post hoc* analysis of the Hypertension in Haemodialysis Patients Treated with Atenolol or Lisinopril (HDPAL) trial [91], each 1-m/s higher baseline aortic PWV was associated with 1.34 mmHg higher 44-h ambulatory SBP and 1.02 mmHg higher PP, but did not predict the treatment-induced reduction in ambulatory SBP and DBP during follow-up. A study evaluating acute changes in arterial stiffness indices during the interdialytic periods showed that augmentation index and central PP increased during both 3- and 2-day interdialytic intervals; aortic and brachial PWV was unchanged in these short time-frames. The increase in augmentation index was 30% greater during the 3-day than during the 2-day interval and was associated with interdialytic weight gain [88]. Subsequent studies with ABPM recordings from the same group further confirmed the above, showing a continuous increase in wave reflection indices and central BP in both the 2- and 3-day interdialytic intervals with minimal increase in PWV [86, 87].

Sympathetic nervous system activation

Seminal microneurography studies suggested that sympathetic overactivity is an important cause of hypertension in ESRD, showing the efferent sympathetic discharge rate to be doubled in haemodialysis patients with *in situ* native kidneys but normal in haemodialysis patients after bilateral nephrectomy [92]. Bilateral nephrectomy of native failed kidneys produced sustained reductions in peripheral vascular resistance and dramatic BP decrease [93]. This pathogenetic role of sympathetic overactivity is also supported by recent observations in which renal denervation substantially reduced BP in small series of haemodialysis patients with severe resistant hypertension [94, 95]. Deficiency of renalase, an enzyme produced by the kidney that metabolizes catecholamines and catecholamine-like substances, may contribute to excessive sympathetic overactivity in CKD [96, 97]. Infusion of recombinant renalase in rats produced a significant reduction in BP, predominantly mediated through reduced peripheral vascular tone and cardiac output [97]. The plasma concentration of renalase is markedly decreased in haemodialysis patients when compared with age- and sex-matched controls with normal renal function [98].

Renin-angiotensin-aldosterone system activation

It is well-known that activation of the renin-angiotensin-aldosterone system occurs even in ESRD patients on renal replacement therapy [99, 100]. Plasma renin activity (PRA) is maintained within the normal range in the majority of patients, but may be inappropriately elevated in relation to the total exchangeable sodium and may contribute to BP elevation [101]. This is supported by clinical studies showing a significant increase in PRA and plasma aldosterone from pre- to

post-dialysis, suggesting that residual functioning nephrons in dialysis patients retain their ability to sense acute changes in sodium intravascular volume status in response to ultrafiltration [99, 101]. Earlier studies showed the angiotensin II antagonist saralasin to lower BP in dialysis [102]; the angiotensin-converting enzyme inhibitor (ACEI) lisinopril was recently shown to reduce 44-h ambulatory BP [103]. The relationship between PRA, aldosterone and major clinical outcomes in dialysis patients is complex and much influenced by malnutrition and inflammation. Independently of pre-dialysis BP, aldosterone is an inverse predictor of cardiovascular events and mortality in this population, and this seemingly paradoxical relationship is abolished by adjustment for inflammation, protein energy malnutrition and volume expansion biomarkers, indicating that it is merely the expression of the confounding effect of these factors [104, 105].

Endothelial dysfunction

An imbalance between endothelium-derived vasodilators and vasoconstrictors may also be involved in hypertension among dialysis patients. Endothelial dysfunction results from several mechanisms. Animal studies document a downregulation of endothelial and inducible NO synthase activity in rats with 5/6 nephrectomy, an alteration that resulted in sustained BP elevation [106]. Patients with CKD also show markedly reduced NO availability, measured as NO-dependent vasodilation [107]. This could be due to reduced production of NO [108], although others describe enhanced NO production in these patients [109]. Increased generation of reactive oxygen species in CKD may cause enhanced breakdown of [110]. Alterations in pteridine metabolism have also been described in chronic renal failure, which may lead to reduced BH4 availability and endothelial NO synthase uncoupling [111]. High circulating levels of asymmetric dimethylarginine (ADMA) [112, 113], an endogenous NO synthase inhibitor, accumulates in CKD and results in reduced generation of NO [114]. The higher levels of ADMA in ESRD result from both diminished intracellular degradation by desamino-D-argininehydrolase and diminished renal clearance of ADMA [114]. Among ESRD patients, ADMA is associated with increased LV relative wall thickness and reduced ejection fraction. Importantly, prospective cohort studies have associated increased ADMA levels with excessive risk of cardiovascular morbidity and mortality in haemodialysis patients [112, 114].

Sleep apnoea

Sleep apnoea is highly prevalent among dialysis patients and volume overload may be a major player in this alteration [115]. In the recumbent position, volume overload may promote sleep-disordered breathing and nocturnal hypoxemia through an overnight fluid shift from the legs to the neck soft tissues that increases peripharyngeal and upper airway resistance [116]. Nocturnal hypoxemia in sleep apnoea has been associated with a reversed circadian BP pattern, in this way triggering nocturnal hypertension. This notion is supported by a study of 32 haemodialysis patients that showed that those patients experiencing sleep apnoea had higher nocturnal SBP and higher LV relative wall thickness than those without sleep apnoea; an inverse relationship was documented between the average nocturnal arterial oxygen saturation and LV relative wall thickness

[35]. In another study, Abdel-Kader *et al.* [117] showed that ESRD patients with sleep apnoea had a 7.1 times higher risk of developing resistant hypertension (defined as office BP >140/90 mmHg despite the use of more than three different antihypertensive agents); in contrast, no such association between sleep apnoea and resistant hypertension was noted among non-dialysis-requiring CKD patients [117]. Finally, a recent study in haemodialysis patients with obstructive sleep apnoea showed that, after haemodialysis, the obstructive apnoea-hypopnoea index was significantly improved only in the group of patients with a concomitant reduction of fluid overload [116]. It remains to be demonstrated whether strict management of volume status restores the blunted nocturnal BP fall in dialysis patients.

Erythropoietin-stimulating agents

Hypertension is a common but frequently overlooked complication of erythropoietin therapy [118]. Hypertension induced by recombinant erythropoietin treatment may depend on increased circulating endothelin-1 or enhanced vasoconstrictive response to endothelin-1 [119, 120], increased sensitivity to the pressor effect of angiotensin II [121], increased blood viscosity and increased vascular sensitivity to noradrenergic stimuli [122]. Higher erythropoietin doses [123], higher target haemoglobin levels [124], route of administration (intravenous versus subcutaneous) [125] and dialysis modality (haemodialysis versus peritoneal dialysis) [126, 127] have all been associated with a higher BP response [128].

Secondary causes of hypertension in dialysis patients

Apart from ESRD and the inability to maintain normal sodium and water homeostasis, practicing nephrologists should not forget that a few patients with hypertension that remain resistant to treatment may have other secondary causes of hypertension, which should be adequately sought and treated [129, 130]. The prevalence and incidence of these disorders can resemble that of the general population, with some exceptions. For example, renovascular disease is rather unlikely to cause hypertension in anuric patients with long dialysis vintage, but should be looked for in patients with heavy atherosclerotic burden, recent dialysis start and residual diuresis. Similarly, primary aldosteronism is unlikely to cause severe hypertension in anuric subjects, as the renal action of aldosterone in maintaining sodium would be absent, but it should be kept in mind for patients with abrupt hypertension and hypokalaemia immediately after kidney transplantation [131]. Obstructive sleep apnoea is particularly common in ESRD patients and is discussed in detail above. Less frequent secondary causes, like pheochromocytoma, thyroid diseases, renin-secreting tumours and others should be carefully sought in selected patients with relevant signs and symptoms, and treated appropriately.

HYPERTENSION TREATMENT IN DIALYSIS PATIENTS

Non-pharmacological measures

Management of hypertension in dialysis patients should focus on correction of the primary pathogenetic mechanism,

that is, sodium and volume excess, by carefully implementing a series of non-pharmacological measures to achieve the dry-weight for each individual patient and to avoid intradialytic sodium loading (Box 3). Particular consideration needs to be given to the fact that, when renal replacement therapy is initiated, 95% of patients are already hypertensive and the vast majority are receiving antihypertensive agents [73]. This, and the fact that common antihypertensive agents may be prescribed for other indications [i.e. β -blockers for angina symptoms, heart failure or rate control, renin-angiotensin system (RAS) blockers for heart failure, etc.], needs to be taken into account and guide the careful handling of antihypertensive drugs when dry-weight is pursued. However, outside situations of hypertensive urgency or emergency [7], the administration of antihypertensive drug therapy in dialysis patients considered to be volume overloaded should follow the attainment of dry-weight.

Achievement of patients' dry-weight. Achievement of dry-weight in dialysis patients remains a complex issue of clinical judgement [132]. The absence of a widely accepted definition of dry-weight and the reliance of definitions on subjective patient symptoms rather than objective estimations are problems known to practicing nephrologists. Sinha and Agarwal [133] defined dry-weight as the lowest tolerated post-dialysis weight, achieved through a gentle and gradual reduction in post-dialysis weight, at which patients experience minimal signs or symptoms of either hypovolaemia or hypervolaemia [133]. Typically, there are no reliable clinical signs to indicate whether a patient has reached the 'ideal' dry-weight. The degree of pedal oedema, which is frequently used as a reference in dialysis patients, was not found to be associated with more objective indices reflecting intravascular volume, such as inferior vena cava diameter, blood volume monitoring or plasma volume biomarkers [134]. In a recent subproject of the ongoing Lung Water by Ultra-Sound Guided Treatment to Prevent Death and Cardiovascular Complications in High Risk ESRD Patients with Cardiomyopathy (LUST) trial, both pedal oedema and crackles in lung auscultation of haemodialysis patients reflected the degree of pulmonary congestion objectively assessed by lung ultrasound very poorly [135]. Bioimpedance methods and relative blood volume monitoring are increasingly used to assess whole body fluid status in dialysis patients [136]; a combination of these methods with lung ultrasound may provide a more precise

Box 3: Main non-pharmacological measures to reduce sodium and volume overload in haemodialysis patients

- Achievement of individual patients' dry-weight.
- Minimization of inter- and intradialytic sodium gain.
 - Restriction of sodium intake to <65 mmol (1.5 g of sodium or 4 g of sodium chloride) per day.
 - Decreasing dialysate sodium towards pre-dialysis sodium in selected individuals.
 - Avoidance of sodium-containing or sodium-exchanging drugs.
- Avoidance of short (i.e. <4 h) dialysis duration.

estimate of fluid accumulation in critical organs and, thus, help towards objective definition of dry-weight [137].

Previous uncontrolled observations in small series of patients [138–140] suggested that supervised gradual reduction (probing) of dry-weight can effectively reduce BP. The DRIP study was the first randomized trial to test this hypothesis, by assigning 150 haemodialysis patients with hypertension in a 2:1 ratio to an intensive ultrafiltration group, in which the dry-weight was probed without increasing the frequency or duration of dialysis, and to a control group, without modification of volume status [141]. In the ultrafiltration group, an initial additional weight loss of 0.1 kg/10 kg body weight was prescribed. If ultrafiltration was not tolerated based on symptoms and signs, such as muscle cramps, a need for excessive saline or symptomatic hypotension, the additional prescribed weight loss was reduced by 50% until 0.2 kg incremental weight loss per dialysis was not tolerated. The primary trial end point was the difference between the ultrafiltration and control groups in the change of 44-h interdialytic ambulatory BP, which was performed at baseline, 4 and 8 weeks. Post-dialysis weight was reduced by 0.9 kg at 4 weeks and resulted in an average difference of 7.4/3.6 mmHg in 44-h interdialytic ambulatory BP between the two groups. The overall dry-weight reduction achieved at study completion was 1 kg and was associated with a difference of 7.1/3.8 mmHg [142]. This benefit was seen without any deterioration in parameters of health-related quality of life [141] and with a parallel reduction in LV chamber volume [142]. Of importance, in the DRIP trial, background antihypertensive treatment of study participants remained unchanged throughout the trial (with an average of 2.7 drugs), indicating that dry-weight reduction can be beneficial even in treated patients. The ongoing LUST trial is a multicentre randomized study within the framework of ERA-EDTA, comparing the effect of dry-weight probing guided by a lung ultrasound scheme versus standard clinical practice on cardiovascular events in haemodialysis patients [143]; a LUST sub-study on ambulatory BP is awaited to shed further light on the field.

In accordance with haemodialysis, achievement of better volume control in patients on peritoneal dialysis may help towards BP normalization. A small, open-label randomized study lasting 12 months showed that, compared with standard glucose peritoneal dialysis solutions, the use of icodextrin solution as an osmotic agent is associated with greater reduction in systolic 24-h ambulatory BP in diabetic patients with high average and high peritoneal transport type [111]. However, in a larger randomized trial comparing a glucose-sparing regimen that included icodextrin with standard glucose peritoneal dialysis solutions in diabetic patients, despite significant improvement in glycated haemoglobin and lipid parameters, deaths and serious adverse events (including those related to volume expansion) increased in the glucose-sparing group [144]. Thus, the optimal way to achieve dry-weight in peritoneal dialysis patients remains to be defined.

Benefits to BP control by an intensification of ultrafiltration in the absence of prolonged dialysis time may be counterbalanced by higher risk of intradialytic hypotension, loss of residual renal function, hospitalization for cardiovascular complications and arteriovenous fistula clotting [5, 145]. High

ultrafiltration rates increase the risk of dialysis hypotension and, in one observational study, ultrafiltration rates >12.4 mL/kg/h were associated with increased mortality [146]. Other uncomfortable symptoms apart from hypotension, such as cramps, nausea and vomiting, may also affect patients' quality of life and interfere with the process of reaching dry-weight. Physicians often respond inappropriately to these symptoms with therapeutic interventions, which may have opposite results to what is intended, such as cessation of ultrafiltration, hypertonic sodium infusions, increasing the dialysate sodium concentration, premature termination of dialysis or finally raising the dry-weight and subsequently increasing the number of prescribed antihypertensive medications (Box 4) [5, 147, 148]. Overall, dry-weight may be more easily and safely achieved in multiple sessions or by prolonging the dialysis time to achieve a slower ultrafiltration rate, as discussed below.

Minimization of inter- and intradialytic sodium gain. As discussed above, in ESRD patients, the sodium and fluid excretory capacity is either absent or substantially impaired and BP is typically salt-sensitive. Thus, reducing the amount of sodium gained from diet or dialysate fluid is critical to achieve BP control. In a cohort study of 1770 haemodialysis patients, high reported dietary sodium (expressed as raw intake, in proportion to caloric intake or in proportion to potassium intake) was associated with greater mortality; of note, adjusted analysis reported that sodium intake displayed a linear association with mortality, starting from the lowest examined levels of 0.5 g/day [149]. Dietary sodium restriction appears to be an effective approach to limit the sense of thirst, reduce interdialytic weight gain and facilitate the achievement of dry-weight and BP control [150]. Observational data suggest that dietary sodium restriction and achievement of dry-weight are associated with improvement of

Box 4: Barriers towards achievement of dry-weight in haemodialysis patients with hypertension

- Difficulty in objectively assessing dry-weight.
- Fear of patient symptoms (intradialytic hypotension, muscle cramps, nausea and vomiting).
- Risk of complications (cardiovascular events, arteriovenous access loss).
- Physician and nurse inertia/ease of prescribing a new drug versus the complex procedure of dry-weight probing.
- Absence of patient education on dietary sodium restriction/misguided emphasis in fluid restriction.
- Low patient compliance with sodium restriction/high interdialytic weight gain.
- Use of sodium-containing medications.
- Inappropriate dialysate sodium.
- Use of high ultrafiltration rates.
- Short dialysis sessions.
- Concomitant diseases (heart failure, autonomic dysfunction).
- Use of high number of antihypertensive agents.
- Use of 'fast and easy' solutions to treat intradialytic hypotension (i.e. cessation of ultrafiltration, hypertonic sodium infusions, increasing dialysate sodium concentration, premature termination of dialysis).

BP and LVH, and fewer episodes of intradialytic hypotension compared with antihypertensive treatment [139, 151]. It should be noted that, in Western countries with frequent consumption of ready meals and processed foods, reducing the amount of sodium intake may be a complex challenge that requires important lifestyle changes. Instead of dietary sodium restriction, patients on dialysis are often instructed to avoid excessive fluid intake during the interdialytic interval; fluid restriction without concomitant sodium restriction is not supported by evidence and is frequently not feasible due to increased thirst [152]. Hypertension guidelines suggest that dietary sodium in any hypertensive patient should be reduced to <100 mmol (2.4 g of sodium or 6 g of sodium chloride) per day [40, 153]. The effect of salt restriction on BP is typically more pronounced in salt-sensitive individuals, like those with CKD; thus, in dialysis patients, dietary sodium intake should not exceed 65 mmol (1.5 g of sodium or 4 g of sodium chloride). In addition, a subset of patients may gain sodium via the use of particular medications, such as potassium-binders exchanging sodium, sodium bicarbonate to increase pre-dialysis bicarbonate levels or drug formulations containing sodium (i.e. effervescent tablets); whenever possible, the avoidance of such agents is also useful.

In parallel to dietary sodium restriction, the avoidance of inappropriate sodium gain during dialysis is crucial for effective BP control. Prescription of a high dialysate sodium concentration was common in the early days of dialysis, to ensure haemodynamic stability and minimize other intradialytic symptoms (i.e. disequilibrium symptoms, nausea, vomiting, muscle cramps, etc.). This was supported by older studies showing that high dialysate sodium may minimize the incidence of intradialytic hypotensive episodes without worsening interdialytic hypertension [154, 155]. However, more recent works have challenged these conclusions and emphasized that a high dialysate sodium concentration may increase thirst and interdialytic weight gain [147, 156]. In a study of 1084 haemodialysis patients, Munoz Mendoza *et al.* [157] found that dialysate sodium prescriptions ranged from 136 to 149 (median, 140) mEq/L and that most patients were dialysed against a positive sodium gradient, resulting in over 90% of patients having a rise in serum sodium across dialysis and, consequently, higher post-dialysis thirst and interdialysis weight gain. Such an increase in interdialytic weight gain leads to a need for greater ultrafiltration during the next dialysis session, which may act as a triggering factor for more frequent episodes of intradialytic hypotension and necessitate the prescription of even higher dialysate sodium concentration, precipitating a vicious cycle [104, 105]. A consensus document by the Chief Medical Officers of US Dialysis Providers warns against the use of dialysate with a sodium concentration exceeding pre-dialysis serum sodium [147, 156].

A single-blind, randomized, cross-over study comparing the effect of nine sessions of a standard dialysate sodium concentration (138 mEq/L) to nine sessions of individualized prescription of the dialysate sodium concentration (the dialysate sodium set to match a patient's average pre-dialysis sodium multiplied by 0.95 to allow for the Gibbs–Donnan effect) in non-diabetic, non-hypotension-prone dialysis patients documented a benefit of individualized sodium prescription on intradialytic weight gain, thirst and episodes of intradialytic BP fall. Among patients

with uncontrolled BP at baseline, pre-dialysis BP was 16 mmHg lower during the individualized sodium dialysate period [158]. In a subsequent single-blind, cross-over study receiving thrice-weekly, in-centre nocturnal dialysis, lowering the dialysate sodium concentration from 140 to 136 or 134 mEq/L for a 12-week treatment period decreased interdialytic weight gain by 0.6 ± 0.6 kg, and pre-dialysis SBP by 8.3 ± 14.9 mmHg, without increasing intradialytic hypotensive episodes [159]. In a 3-week randomized, cross-over trial in 16 patients with intradialytic hypertension, Inrig *et al.* [50] compared the effect of a high (5 mEq/L above serum sodium) versus low (5 mEq/L below serum sodium) dialysate sodium concentration on intradialytic BP and endothelial-derived vasoregulators. The weekly averaged pre-dialysis SBP was lower during the period of low dialysate sodium concentration (-9.9 mmHg; 95% CI: -13.3 to -6.4 mmHg; $P < 0.001$), as was the weekly average intradialytic SBP (-6.1 mmHg; 95% CI: -9.0 to -3.2 mmHg; $P < 0.001$) (Figure 3) [50]. Overall, these studies suggest that a single dialysate sodium prescription may not fit all patients. Small decreases in dialysate sodium towards pre-dialysis levels in hypertensive patients can limit thirst, reduce intradialytic weight gain and improve BP control without aggravating the risk of intradialytic haemodynamic instability. Larger randomized trials are needed to evaluate the safety and efficacy of this approach.

In peritoneal dialysis patients, increasing the diffusive component of sodium removal with the use of low-sodium peritoneal dialysis fluids is suggested to be an effective intervention to improve BP control. In a non-randomized interventional study comparing a standard versus a low-sodium peritoneal dialysis solution substituting one 3- to 5-h exchange per day over a mean follow-up period of 2 months, the use of low-sodium dialysate resulted in a significant increase of 30–50 mmol/dwell diffusive peritoneal sodium removal, which was accompanied by reduced thirst, lower total body water and an 8 mmHg fall of night-time SBP [160]. Overall, patients on peritoneal dialysis should also follow the above recommendations for restriction of sodium intake; modification of peritoneal dialysis regimens with low-sodium or icodextrin solutions may facilitate sodium and volume control.




FIGURE 3: Weekly average intradialytic systolic blood pressure (SBP) during haemodialysis with low and high dialysate sodium (Na) by randomization sequence (group 1: low-then-high dialysate Na) and (group 2: high-then-low dialysate Na); $P < 0.001$ by adjusted mixed linear regression. Reprinted with permission from Inrig *et al.* [50].

Avoiding short dialysis. Among several other potential hazards, short delivered dialysis can be an important barrier to the achievement of adequate BP control. The European Best Practice guidelines recommend that the length of the dialysis session must not be decided only on the grounds of optimal Kt/V and that haemodialysis patients should receive at least three dialysis sessions of 4 h each per week [161], a recommendation aimed mainly at ensuring optimal volume status. Exceptions to this could be incident dialysis patients with substantial residual renal function or patients who start dialysis early during the evolution of their CKD; these specific subgroups of dialysis patients may be able to maintain the homeostasis of volume and metabolic parameters over a longer dialysis-free interval [161–164]. However, real world data deriving from registries throughout the globe suggest that the reality is different and that, although the mean dialysis session length may be around 210–235 min, some patients may receive dialysis for shorter times; this is particularly relevant for registries in the USA, where as many as 25% of patients may dialyse for <3 h and 15 min per session [165–167].

Increasing the duration of dialysis may represent an additional approach to control BP among dialysis patients who remain hypertensive despite the intensification of volume withdrawal, or who experience frequent episodes of intradialytic haemodynamic instability during this intensification process within their usual dialysis regimen [168]. A previous cross-over study of 38 dialysis patients comparing the frequency of intradialytic symptoms during 5-h versus 4-h dialysis sessions showed that the incidence of intradialytic hypotension and post-dialysis orthostatic hypotension was less common during the period of extended-time dialysis [169]. This notion is supported by a *post hoc* analysis of the DRIP trial [126], in which median intradialytic SBP at baseline and its change over time were modelled against the duration of delivered dialysis. At baseline, median intradialytic SBP was higher with fewer hours of delivered dialysis. Among patients in whom dry-weight was not reduced (control group), median intradialytic SBP followed an increasing trend over the course of the trial. In the ultrafiltration group, dry-weight reduction induced a significant drop in median intradialytic SBP regardless of the duration of delivered dialysis. However, patients with longer delivered dialysis required fewer dialysis sessions in order to gain the BP-lowering benefit of dry-weight reduction. A similar beneficial relationship was evident between the duration of delivered dialysis and the magnitude of change in 44-h interdialytic ambulatory SBP over time [126].

The fact that avoiding short dialysis may facilitate BP control is also supported by several other randomized and non-randomized studies, showing that patients assigned to longer (i.e. up to 8-h thrice weekly) or more frequent (i.e. up to six times/week) dialysis regimens achieve better BP control with reduced requirements for antihypertensive medications. This benefit is possibly mediated through better correction of sodium and volume excess [168, 170–172].

Of note, a recent long-term, observational post-trial analysis of patients who took part into the Daily in-centre Trial of the Frequent Haemodialysis Network [173] showed a lower risk of death in patients originally randomized to frequent haemodialysis (six times a week) and 1.5–2.75 h/session (16%), as

compared with those randomized to conventional haemodialysis treatment (28%). However, this benefit was not evident in the long-term analysis of the twin Nocturnal Trial of the same network, where mortality was largely increased in the frequent haemodialysis group (six times a week >6 h/session) [174]. Of note, the most prominent difference between groups in the main Nocturnal Trial seemed to be a faster loss of residual diuresis in the frequent dialysis arm [175]. Although careful interpretation is necessary, current evidence suggests that longer or frequent haemodialysis schemes may be beneficial, but that the combination of both longer and frequent treatment is not.

Pharmacological treatment

The effects of β -blockers, ACEIs, angiotensin-II receptor blockers (ARBs), calcium channel blockers (CCBs) and mineralocorticoid receptor antagonists (MRAs) on hard outcomes in haemodialysis patients have been examined in clinical trials (Box 5). Two previous meta-analyses of randomized trials clearly suggest that BP-lowering with the use of such

Box 5: Antihypertensive drugs in outcome clinical trials in haemodialysis patients

β -blockers

- Carvedilol reduced mortality compared with placebo in haemodialysis patients with dilated cardiomyopathy [176].
- Thrice-weekly atenolol reduced cardiovascular events compared with thrice-weekly lisinopril in HD patients with hypertension and LVH in the HDPAL trial [177].

ACEIs

- Fosinopril did not reduce cardiovascular events and mortality compared with placebo in HD patients with LVH in the FOSIDIAL trial [178].

ARBs

- Losartan/valsartan/candesartan reduced cardiovascular events and mortality compared with treatment not including ACEIs/ARBs in HD patients [179, 180].
- Olmesartan did not reduce cardiovascular events or mortality compared with treatment not including ACEIs/ARBs in HD patients with hypertension in the OCTOPUS trial [181].

CCBs

- Amlodipine reduced cardiovascular events compared with placebo in HD patients with hypertension [182].

MRAs

- Spironolactone may reduce cardiovascular events and mortality compared with no additional treatment or placebo in HD and peritoneal patients [183, 184].

HD, haemodialysis; LVH, left ventricular hypertrophy; HDPAL, Hypertension in Haemodialysis Patients Treated with Atenolol or Lisinopril trial; ACEIs, angiotensin-converting enzyme inhibitors; FOSIDIAL, Fosinopril in Dialysis trial; ARBs, angiotensin-II receptor blockers; OCTOPUS, Olmesartan Clinical Trial in Okinawa Patients under Dialysis Study; CCBs, calcium channel blockers; MRAs, mineralocorticoid receptor antagonists.

antihypertensive drugs is associated with reduced cardiovascular morbidity and mortality in dialysis patients [185, 186]. The first meta-analysis included eight trials incorporating data from 1697 dialysis patients and 495 cardiovascular events [186]. The weighted mean difference in the change of BP between the active treatment and control groups was -4.5 mmHg for SBP and -2.3 mmHg for DBP. This BP-lowering effect of antihypertensive drug treatment was associated with a 29% reduction in the risk of all-cause mortality (pooled RR: 0.71; 95% CIs: 0.55–0.92) and a 29% reduction in the risk of cardiovascular mortality (pooled RR: 0.71; 95% CIs: 0.50–0.99) [186]. The second meta-analysis [185] included five randomized trials with 1202 study participants. Compared with placebo or control therapy, the overall cardiovascular benefit of BP-lowering with antihypertensive therapy was a 31% reduction in the risk of future cardiovascular events (pooled HR: 0.69; 95% CIs: 0.56–0.84) [185]. In a subanalysis according to the hypertension status of patients participating in the individual studies, it was shown that cardiovascular protection provided by BP-lowering was less pronounced when normotensive patients were included in the analysis (pooled HR: 0.86; 95% CIs: 0.67–1.12) [185]. These meta-analyses indicate that the use of antihypertensive drugs in dialysis patients may afford cardiovascular protection both in hypertensive patients and in normotensive patients with LV systolic dysfunction [185].

The major antihypertensive drug classes are useful for the pharmacological treatment of hypertension in dialysis, taking into account the specific pharmacological properties of each drug [5, 9, 187, 188]. An exception may be diuretics, which are ineffective for BP control in patients with ESRD [5, 187, 188]. Echocardiographic studies conducted in anuric haemodialysis patients showed that intravenous administration of loop diuretics, even at high doses, exerts only minimal alterations in central haemodynamic indices [189]. Given the high risk of ototoxicity, the use of loop diuretics in anuric dialysis patients should be avoided. Several small studies suggest that these compounds may help patients with preserved residual diuresis on haemodialysis or peritoneal dialysis to enhance urine output and limit fluid overload [190–193]; however, the effect of loop diuretics on urine output and BP control has not been properly examined in large studies.

β -blockers. Sympathetic overactivity, as measured by plasma norepinephrine, is a powerful predictor of death and cardiovascular events in dialysis patients [194]. The susceptibility of dialysis patients to serious arrhythmias and sudden death, along with the excessive activation of the sympathetic nervous system, make β -blockers an attractive therapeutic option for cardiovascular protection in this population [187]. Interestingly, in an analysis of the DOPPS study, use of β -blockers was associated with a lower risk of sudden death, after adjustment for comorbidities (HR: 0.88; 95% CI: 0.78–0.99; $P=0.03$) [195]. In 114 haemodialysis patients with dilated cardiomyopathy randomized to carvedilol (up to 25 mg twice daily) or placebo for 2 years, carvedilol improved LV systolic function and significantly reduced the risk of all-cause hospitalization (HR: 0.44; 95% CI: 0.25–0.77) and all-cause mortality (HR: 0.51; 95% CI: 0.32–0.82) [176]. More recently, the HDPAL trial [177] performed a

head-to-head comparison between the β -blocker atenolol and the ACEI lisinopril (both administered in a thrice-weekly regimen immediately post-dialysis) in 200 hypertensive haemodialysis patients with echocardiographically documented LVH. The trial showed that the LV mass index over the 12-month follow-up (the primary outcome) improved to a similar extent in the atenolol and lisinopril groups [177]. However, atenolol was shown to be superior to lisinopril in terms of its BP-lowering efficacy; in particular, no significant differences in BP were noted between the two groups, but lisinopril-treated patients had always numerically higher BP levels (Figure 4), required more aggressive volume management during dialysis and the administration of a higher number of antihypertensive drugs as add-on therapy to achieve the pre-specified home BP target of 140/90 mmHg. Most importantly, the HDPAL trial was terminated early due to the superiority of atenolol over lisinopril for the prevention of serious cardiovascular events, as the rate of the combined outcome of myocardial infarction (MI), stroke, hospitalized heart failure and cardiovascular death was 2.29 times higher in lisinopril-treated than atenolol-treated patients (incidence rate ratio: 2.29; 95% CI: 1.07–5.21) [177].

The Beta-blocker to LOwer Cardiovascular Dialysis Events trial originally planned to study the cardioprotective role of β -blockade in haemodialysis patients. In the feasibility study, which aimed to enrol 150 patients, among the 1443 patients screened (including 176 who were already on treatment with β -blockers), only 354 were eligible, 91 consented and 72 entered the 6-week active treatment run-in period. Of these, only 49 participants (68%; 95% CI: 57–79%) tolerated carvedilol therapy (6.25 mg twice daily) during the run-in period and progressed to randomization [196]. The challenging recruitment for this study emphasizes the difficulties in performing clinical studies in dialysis patients.

Pilot data by Inrig *et al.* [197] suggest that carvedilol may be useful in patients with intradialytic hypertension; the authors showed that carvedilol treatment in these patients was associated with an improvement in endothelium-dependent flow-mediated vasodilatation. This effect was accompanied by reduced occurrence of intradialytic hypertensive episodes during follow-up and a significant drop of 7 mmHg in 44-h interdialytic ambulatory SBP. Of importance, when prescribing a β -blocker to a haemodialysis patient, one needs to take into account that there are major differences in renal clearance and dialysability between different agents of this class, as discussed in detail elsewhere [6]. Use of non-dialysable β -blockers is advisable, since a recent retrospective cohort study suggested that a survival advantage may not be offered by highly dialysable β -blockers, possibly due to a lack of intradialytic protection against arrhythmias as a consequence of rapid removal with dialysis [198].

Angiotensin-converting enzyme inhibitor and angiotensin receptor blockers. Blockers of the RAS are among the most widely used antihypertensive agents worldwide. Of note, ACEIs and ARBs are not interchangeable for dialysis patients, as there are important differences between in their renal clearance and removal during dialysis [6, 9]; most ARBs are not dialysed during conventional dialysis and may be preferred in these patients




FIGURE 4: Blood pressure measured by 44-h ABPM over the interdialytic period (left panel) and self-measured by the patients at home (right panel) in the Hypertension in Hemodialysis Patients Treated with Atenolol or Lisinopril (HDPAL) trial. Dotted lines: lisinopril group; solid lines: atenolol group. Reprinted with permission from Agarwal *et al.* [177]. Solid line shows the atenolol group and the dotted line the lisinopril group; vertical bars represent standard error of mean. The table at the bottom of each graph shows the number of patients in each drug [atenolol (n), lisinopril (n)]; the change from baseline (CFB) and between group comparisons of the changes (lisinopril-atenolol CFB).

for sustained BP reduction. Through extrapolation of the cardiovascular benefits of RAS blockers in the general population, inhibition of the RAS has often been recommended as the first-line BP-lowering therapy for dialysis patients [40]. However, randomized trials in hypertensive dialysis patients do not support the notion that RAS blockade offers the same benefits as in hypertensive patients in the general population.

In the Fosinopril in Dialysis trial [178], 397 haemodialysis patients were randomized to receive the ACEI fosinopril (titrated up to 20 mg/day) or placebo for a mean follow-up period of 48 months. Participating patients had per protocol LVH, but were not necessarily hypertensive. Although therapy with fosinopril resulted in a significant reduction of pre-dialysis BP versus placebo in the subgroup of hypertensive participants, the occurrence of fatal and non-fatal cardiovascular events during the follow-up period did not differ significantly between the active treatment and placebo arms (RR: 0.93; 95% CI: 0.68–1.26) [178].

Three trials [179–181], all performed in Japan, compared ARBs to either placebo or active therapy. In two of these trials (including 80 and 360 haemodialysis patients, respectively) the risk of cardiovascular events was remarkably lower in patients treated with ARBs. In the third study, the Olmesartan Clinical Trial in Okinawa Patients under Dialysis Study [181], which was also the largest to date, 469 hypertensive haemodialysis patients were randomized to the ARB olmesartan (10–40 mg/day)

or control therapy not including ACEIs or ARBs. Over a mean follow-up of 3.5 years, and for similar BP control, incidence of all-cause death, non-fatal stroke, MI and coronary revascularization was similar in the olmesartan and control groups (HR: 1.00; 95% CI: 0.71–1.40) [181], suggesting that antihypertensive treatment *per se*, and not the use of an RAS blocker, is the factor responsible for reducing cardiovascular risk. A meta-analytical estimate of the risk reduction by ARBs in these trials (which included around 900 patients and 175 deaths) showed a non-significant ($P = 0.10$) 42% risk reduction [199]. Overall, to date, superiority of ACEIs and ARBs over other antihypertensive drugs has not been demonstrated in dialysis patients, and antihypertensive treatment *per se* rather than the use of an RAS blocker seems to be the factor reducing cardiovascular risk.

Calcium channel blockers. Dihydropyridine CCBs are potent antihypertensive agents that can effectively lower BP, even in the volume-expanded state [200], and are often used for the management of hypertension in dialysis patients. In the only relevant study examining hard outcomes, Tepel *et al.* [182] randomized 251 hypertensive haemodialysis patients to receive amlodipine (5–10 mg/day) or placebo for 30 months. Amlodipine improved survival compared with placebo, although this was not significant, and reduced by 47% the composite secondary end point of all-cause death, non-fatal stroke, MI, coronary revascularization and angioplasty for peripheral

vascular disease (HR: 0.53; 95% CI: 0.31–0.93) [182]. Small previous studies have suggested that dihydropyridine CCBs are equally effective as ACEIs or ARBs in reducing LVH and carotid intima-media thickness [201]. Data on non-dihydropyridine CCB use in haemodialysis patients are scarce; using these agents should at least follow the recommendations for the general population. It must be noted that all CCBs are not removed during standard haemodialysis and their pharmacokinetics are unchanged in ESRD; thus, they can be dosed once-daily in these patients [6, 9].

Mineralocorticoid receptor antagonists. A cardioprotective action of MRAs in dialysis patients has solid biological underpinnings [202], and two recent trials (Table 2) [183, 184] apparently support the contention that these drugs may provide substantial benefits in dialysis patients. In the Dialysis Outcomes Heart Failure Aldactone Study, 309 oligoanuric haemodialysis patients were randomized to spironolactone (25 mg/day) or no add-on therapy for 3 years. Spironolactone reduced the risk of cardiovascular mortality or cardiovascular-related hospitalization (HR: 0.38; 95% CI: 0.17–0.83), with the incidence of drug discontinuation due to serious hyperkalaemia being 1.9% and due to adverse effects overall being 14.6% [183]. In another study, 253 haemodialysis or peritoneal dialysis patients without heart failure were randomized to 2-year-long add-on therapy with spironolactone (25 mg/day) or placebo. Add-on MRA therapy again reduced the occurrence of the composite primary end point of cardio-cerebrovascular mortality, and mitigated the risk for cardiac arrest and sudden death (HR: 0.42; 95% CI: 0.26–0.78) [184]. The reduction in the risk of adverse clinical outcomes in these trials exceeded 50%, that is, it was apparently superior to the effect of frequent in-centre haemodialysis on the combined end point of death and LVH progression [170]; this was largely unexpected in a population like the ESRD population, which is notoriously less sensitive to interventions aimed at reducing death and cardiovascular events than other patient populations [203]. However, it has to be noted that these results should be further confirmed, as both of the above studies were open-label. The safety profile of MRAs in the dialysis population was investigated in a recent study, in which 146 haemodialysis patients were randomly assigned to eplerenone (25–50 mg daily) or matching placebo for 13 weeks [204]. Eplerenone treatment significantly increased the incidence of hyperkalaemia (defined as pre-dialysis serum potassium >6.5 mmol/L) as compared with placebo (RR: 4.50; 95% CI: 1.0–20.2) [204], but permanent drug discontinuation due to hyperkalaemia or hypotension, which was the primary study end point, was no different between eplerenone and placebo groups [204]. Adequately powered, properly designed studies, like the ongoing [205] ALdosterone Antagonist Chronic HEModialysis Interventional Survival Trial (NCT01848639) are needed to assess the effectiveness and safety of mineralocorticoid receptor blockade in ESRD, prior to the recommendation of wider use of MRAs in this population.

Table 2. Recent randomized trials on the effect of MRAs on cardiovascular outcomes in haemodialysis patients

Author	Patient characteristics	N	Design	Follow-up	BP medication	BP assessment	Baseline BP (mmHg)	Final BP (mmHg)	Main finding
Matsumoto <i>et al.</i> [183]	Oligoanuric HD patients	157 versus 152	Open-label RCT	36 months	Spironolactone versus nothing	Pre-dialysis BP	152.8/77.8 versus 148.8/76.2	152.7/77.9 versus N/A	Spironolactone reduced the risk of death or hospitalization for CV event (HR: 0.38; 95% CI: 0.17–0.83)
Lin <i>et al.</i> [184]	HD or PD patients without CHF	125 versus 128	Open-label RCT	24 months	Spironolactone versus placebo	Pre-dialysis BP	144.7/76.9 versus 141.9/77.4	N/A	Spironolactone reduced the risk of CV death, sudden death or aborted cardiac arrest (HR: 0.42; 95% CI: 0.26–0.78)

MRAs, mineralocorticoid receptor antagonists; BP, blood pressure; CV, cardiovascular; CHF, congestive heart failure; CI, confidence intervals; HD, haemodialysis; HR, hazard ratio; LVH, left ventricular hypertrophy; PD, peritoneal dialysis; RCT, randomized clinical trial; N/A, not applicable.

CONCLUSION

Hypertension in dialysis patients poses almost unique diagnostic, prognostic and therapeutic challenges. The evolution of studies using home or ambulatory BP monitoring is currently needed in order to better define the true burden of hypertension in haemodialysis and peritoneal dialysis patients, to provide solid data on hypertension prevalence and prognostic associations and to identify objective thresholds for diagnosis and targets for treatment. Non-pharmacological interventions targeting sodium and volume excess are fundamental for BP reduction in this population and should be carefully implemented before pharmacological interventions. Among dialysis patients, BP-lowering with the use of antihypertensive agents is associated with improvement in cardiovascular outcomes; the use of β -blockers followed by dihydropyridine CCBs should be considered. The first-line use of ACEIs and ARBs in this population

Box 6: Areas in the field of hypertension in dialysis patients where future research efforts are needed

Epidemiology

- Validation studies of devices used for BP recording during dialysis.
- Studies testing the applicability and tolerance of ambulatory BP monitoring and the availability of patients for repeated measurements over time.
- Studies using home or ambulatory BP monitoring to define the true burden of hypertension in HD and PD patients.
- Comparative studies using office, home and ambulatory BP monitoring to further delineate their predictive power for cardiovascular events and death.
- Randomized clinical trials with different BP targets to objectively identify targets for treatment.

Pathophysiology

- Human studies to delineate the interplay between established mechanisms (e.g. between variations on volume and sodium load and changes in other mechanisms) and to uncover novel pathogenic pathways.
- Studies to define novel, objective tools to measure volume overload.

Treatment

- Further clinical trials on the effect of non-pharmacologic interventions (i.e. dry-weight reduction based on objective tools—e.g. the LUST study [143]—restriction of dietary sodium based on objective dietary instruments, increased duration of dialysis, etc.) on home or ambulatory BP control and hard outcomes.
- Further clinical trials on the effect of pharmacologic interventions (i.e. a head-to-head comparison of everyday use of β -blockers versus ACEI/ARBs or CCBs, a proper placebo-controlled trial with an MRA, etc.) on home or ambulatory BP control and hard outcomes.

BP, blood pressure; HD, haemodialysis; PD, peritoneal dialysis; LUST, Lung Water by Ultra-Sound Guided Treatment to Prevent Death and Cardiovascular Complications in High Risk ESRD Patients with Cardiomyopathy study; ACEIs, angiotensin-converting enzyme inhibitors; ARBs, angiotensin-II receptor blockers; CCBs, calcium channel blockers; MRA, mineralocorticoid receptor antagonist.

is not supported by randomized trials. Furthermore, properly designed epidemiology studies and clinical trials to define BP targets for treatment and examine the efficacy of non-pharmacologic measures in reducing BP and antihypertensive drugs in the prevention of major cardiovascular outcomes in the ESRD population remain a public health priority (Box 6).

Alberto Ortiz was supported by FEDER and Spanish government funds ISCIII RETIC REDNREN RD012/0021 and Intensificacion.

CONFLICT OF INTEREST STATEMENT

R.A. has consulted for Abbvie, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Celgene, Daiichi Sankyo Inc., Eli Lilly, Gilead, GlaxoSmithKline, Johnson & Johnson, Merck, Novartis, Sandoz, Relypsa and ZS Pharma. The remaining authors report no conflict of interest relevant to this work.

ACKNOWLEDGEMENTS

Alberto Ortiz was supported by European Regional Development Fund (ERDF FEDER) and Spanish government funds Instituto de Salud Carlos III Red Temática de Investigación Clínica (ISCIII RETIC) Red de Investigación Renal (REDINREN) RD012/0021, RD016/0009 and Intensificacion. The authors wish to thank Drs P. Georgianos and C. Loutradis for updating the literature search relevant to this manuscript.

REFERENCES

1. Agarwal R, Nissenson AR, Battle D *et al.* Prevalence, treatment, and control of hypertension in chronic hemodialysis patients in the United States. *Am J Med* 2003; 115: 291–297
2. Agarwal R. Blood pressure and mortality among hemodialysis patients. *Hypertension* 2010; 55: 762–768
3. Alborzi P, Patel N, Agarwal R. Home blood pressures are of greater prognostic value than hemodialysis unit recordings. *Clin J Am Soc Nephrol* 2007; 2: 1228–1234
4. Amar J, Vernier I, Rossignol E *et al.* Nocturnal blood pressure and 24-hour pulse pressure are potent indicators of mortality in hemodialysis patients. *Kidney Int* 2000; 57: 2485–2491
5. Agarwal R, Flynn J, Pogue V *et al.* Assessment and management of hypertension in patients on dialysis. *J Am Soc Nephrol* 2014; 25: 1630–1646
6. Levin NW, Kotanko P, Eckardt KU *et al.* Blood pressure in chronic kidney disease stage 5D-report from a Kidney Disease: Improving Global Outcomes controversies conference. *Kidney Int* 2010; 77: 273–284
7. Zoccali C, Mallamaci F. Hypertension. In: JT Daugirdas, PG Blake, TS Ing, eds. *Handbook of Dialysis*. 5th edn. Philadelphia: Wolters Kluwer, 2015, 578–591
8. Agarwal R, Martinez-Castelao A, Wiecek A *et al.* The lingering dilemma of arterial pressure in CKD: what do we know, where do we go? *Kidney Int Suppl* 2011; 1: 17–20
9. Redon J, Martinez F, Cheung AK. Special considerations for antihypertensive agents in dialysis patients. *Blood Purif* 2010; 29: 93–98
10. K/DOQI clinical practice guidelines on hypertension and antihypertensive agents in chronic kidney disease. *Am J Kidney Dis* 2004; 43: S1–S290
11. Parati G, Ochoa JE, Bilo G *et al.* Hypertension in chronic kidney disease part 1: out-of-office blood pressure monitoring: methods, thresholds, and patterns. *Hypertension* 2016; 67: 1093–1101

12. Parati G, Ochoa JE, Bilo G *et al.* Hypertension in chronic kidney disease part 2: role of ambulatory and home blood pressure monitoring for assessing alterations in blood pressure variability and blood pressure profiles. *Hypertension* 2016; 67: 1102–1110
13. Rahman M, Griffin V, Kumar A *et al.* A comparison of standardized versus “usual” blood pressure measurements in hemodialysis patients. *Am J Kidney Dis* 2002; 39: 1226–1230
14. Rohrscheib MR, Myers OB, Servilla KS *et al.* Age-related blood pressure patterns and blood pressure variability among hemodialysis patients. *Clin J Am Soc Nephrol* 2008; 3: 1407–1414
15. Agarwal R. Volume-associated ambulatory blood pressure patterns in hemodialysis patients. *Hypertension* 2009; 54: 241–247
16. Inrig JK, Patel UD, Gillespie BS *et al.* Relationship between interdialytic weight gain and blood pressure among prevalent hemodialysis patients. *Am J Kidney Dis* 2007; 50: 108–118
17. Agarwal R, Peixoto AJ, Santos SF *et al.* Pre- and postdialysis blood pressures are imprecise estimates of interdialytic ambulatory blood pressure. *Clin J Am Soc Nephrol* 2006; 1: 389–398
18. Agarwal R, Andersen MJ, Bishu K *et al.* Home blood pressure monitoring improves the diagnosis of hypertension in hemodialysis patients. *Kidney Int* 2006; 69: 900–906
19. Zoccali C, Tripepi R, Torino C *et al.* Moderator’s view: ambulatory blood pressure monitoring and home blood pressure for the prognosis, diagnosis and treatment of hypertension in dialysis patients. *Nephrol Dial Transplant* 2015; 30: 1443–1448
20. Agarwal R, Satyan S, Alborzi P *et al.* Home blood pressure measurements for managing hypertension in hemodialysis patients. *Am J Nephrol* 2009; 30: 126–134
21. Gorostidi M, Sarafidis PA, de la Sierra A *et al.* Differences between office and 24-hour blood pressure control in hypertensive patients with CKD: A 5,693-patient cross-sectional analysis from Spain. *Am J Kidney Dis* 2013; 62: 285–294
22. Mnif K, Jarraya F, Chaker H *et al.* [PP.28.14] Benefit of forty-eight-hour ambulatory blood pressure monitoring in hemodialysis population. *J Hypertens* 2016; 34: e298
23. Agarwal R, Metiku T, Teegne GG *et al.* Diagnosing hypertension by intradialytic blood pressure recordings. *Clin J Am Soc Nephrol* 2008; 3: 1364–1372
24. Parati G, Stergiou GS, Asmar R *et al.* European Society of Hypertension guidelines for blood pressure monitoring at home: a summary report of the Second International Consensus Conference on Home Blood Pressure Monitoring. *J Hypertens* 2008; 26: 1505–1526
25. Agarwal R, Light RP. Median intradialytic blood pressure can track changes evoked by probing dry-weight. *Clin J Am Soc Nephrol* 2010; 5: 897–904
26. Agarwal R, Brim NJ, Mahenthiran J *et al.* Out-of-hemodialysis-unit blood pressure is a superior determinant of left ventricular hypertrophy. *Hypertension* 2006; 47: 62–68
27. Moriya H, Ohtake T, Kobayashi S. Aortic stiffness, left ventricular hypertrophy and weekly averaged blood pressure (WAB) in patients on haemodialysis. *Nephrol Dial Transplant* 2007; 22: 1198–1204
28. Moriya H, Oka M, Maesato K *et al.* Weekly averaged blood pressure is more important than a single-point blood pressure measurement in the risk stratification of dialysis patients. *Clin J Am Soc Nephrol* 2008; 3: 416–422
29. Agarwal R, Andersen MJ, Light RP. Location not quantity of blood pressure measurements predicts mortality in hemodialysis patients. *Am J Nephrol* 2008; 28: 210–217
30. Viazzi F, Leoncini G, Ratto E *et al.* Peripheral artery disease and blood pressure profile abnormalities in hemodialysis patients. *J Nephrol* 2016 [epub ahead of print]
31. da Silva GV, de BS, Abensur H *et al.* Home blood pressure monitoring in blood pressure control among haemodialysis patients: an open randomized clinical trial. *Nephrol Dial Transplant* 2009; 24: 3805–3811
32. Kauric-Klein Z, Artinian N. Improving blood pressure control in hypertensive hemodialysis patients. *CANNT J* 2007; 17: 24–26
33. Agarwal R. Pro: Ambulatory blood pressure should be used in all patients on hemodialysis. *Nephrol Dial Transplant* 2015; 30: 1432–1437
34. Sarafidis PA, Rumjon A, Macdougall IC. Ambulatory blood pressure monitoring: an invaluable tool comes of age for patients with chronic kidney disease? *Am J Nephrol* 2012; 35: 238–241
35. Zoccali C, Benedetto FA, Tripepi G *et al.* Nocturnal hypoxemia, night-day arterial pressure changes and left ventricular geometry in dialysis patients. *Kidney Int* 1998; 53: 1078–1084
36. Tripepi G, Fagugli RM, Dattolo P *et al.* Prognostic value of 24-hour ambulatory blood pressure monitoring and of night/day ratio in nondiabetic, cardiovascular events-free hemodialysis patients. *Kidney Int* 2005; 68: 1294–1302
37. O’Brien E, Parati G, Stergiou G *et al.* European Society of Hypertension position paper on ambulatory blood pressure monitoring. *J Hypertens* 2013; 31: 1731–1768
38. Hypertension: clinical management of primary hypertension in adults. *NICE Clinical Guideline* 127 2011, <http://guidance.nice.org.uk/CG127> (2 November 2016, date last accessed)
39. Piper MA, Evans CV, Burda BU *et al.* Diagnostic and predictive accuracy of blood pressure screening methods with consideration of rescreening intervals: a systematic review for the U.S. preventive services task force-blood pressure screening methods and consideration of rescreening intervals. *Ann Int Med* 2015; 162: 192–204
40. Mancia G, Fagard R, Narkiewicz K *et al.* 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013; 31: 1281–1357
41. Parati G, Stergiou G, O’Brien E *et al.* European Society of Hypertension practice guidelines for ambulatory blood pressure monitoring. *J Hypertens* 2014; 32: 1359–1366
42. Bansal N, McCulloch CE, Rahman M *et al.* Blood pressure and risk of all-cause mortality in advanced chronic kidney disease and hemodialysis: the chronic renal insufficiency cohort study. *Hypertension* 2015; 65: 93–100
43. Sarafidis PA, Ruilope LM. Aggressive blood pressure reduction and renin-angiotensin system blockade in chronic kidney disease: time for re-evaluation? *Kidney Int* 2014; 85: 536–546
44. Wright JT Jr, Williamson JD, Whelton PK *et al.* A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med* 2015; 373: 2103–2116
45. Jardine AG, Agarwal R. Con: Ambulatory blood pressure measurement in patients receiving haemodialysis: a sore arm and a waste of time? *Nephrol Dial Transplant* 2015; 30: 1438–1441
46. Georgianos PI, Sarafidis PA, Zoccali C. Intradialysis hypertension in end-stage renal disease patients: clinical epidemiology, pathogenesis, and treatment. *Hypertension* 2015; 66: 456–463
47. Inrig JK. Intradialytic hypertension: a less-recognized cardiovascular complication of hemodialysis. *Am J Kidney Dis* 2010; 55: 580–589
48. Raj DS, Vincent B, Simpson K *et al.* Hemodynamic changes during hemodialysis: role of nitric oxide and endothelin. *Kidney Int* 2002; 61: 697–704
49. Inrig JK, Patel UD, Toto RD *et al.* Association of blood pressure increases during hemodialysis with 2-year mortality in incident hemodialysis patients: a secondary analysis of the Dialysis Morbidity and Mortality Wave 2 Study. *Am J Kidney Dis* 2009; 54: 881–890
50. Inrig JK, Molina C, D’Silva K *et al.* Effect of low versus high dialysate sodium concentration on blood pressure and endothelial-derived vasoregulators during hemodialysis: a randomized crossover study. *Am J Kidney Dis* 2014; 65: 464–473
51. Agarwal R, Light RP. Intradialytic hypertension is a marker of volume excess. *Nephrol Dial Transplant* 2010; 25: 3355–3361
52. Van Buren PN, Kim C, Toto R *et al.* Intradialytic hypertension and the association with interdialytic ambulatory blood pressure. *Clin J Am Soc Nephrol* 2011; 6: 1684–1691
53. Rahman M, Fu P, Sehgal AR *et al.* Interdialytic weight gain, compliance with dialysis regimen, and age are independent predictors of blood pressure in hemodialysis patients. *Am J Kidney Dis* 2000; 35: 257–265
54. Rocco MV, Yan G, Heyka RJ *et al.* Risk factors for hypertension in chronic hemodialysis patients: baseline data from the HEMO study. *Am J Nephrol* 2001; 21: 280–288
55. Salem MM. Hypertension in the hemodialysis population: a survey of 649 patients. *Am J Kidney Dis* 1995; 26: 461–468
56. Agarwal R. Epidemiology of interdialytic ambulatory hypertension and the role of volume excess. *Am J Nephrol* 2011; 34: 381–390

57. Sarafidis PA, Li S, Chen SC *et al.* Hypertension awareness, treatment, and control in chronic kidney disease. *Am J Med* 2008; 121: 332–340
58. Sarafidis PA, Sharpe CC, Wood E *et al.* Prevalence, patterns of treatment, and control of hypertension in predialysis patients with chronic kidney disease. *Nephron Clin Pract* 2012; 120: c147–c155
59. Katzarski KS, Charra B, Luik AJ *et al.* Fluid state and blood pressure control in patients treated with long and short haemodialysis. *Nephrol Dial Transplant* 1999; 14: 369–375
60. Rahman M, Dixit A, Donley V *et al.* Factors associated with inadequate blood pressure control in hypertensive hemodialysis patients. *Am J Kidney Dis* 1999; 33: 498–506
61. Hypertension, by DOPPS Country and Cross-Section. 2006. <http://www.dopps.org/annualreport/html/MCOMHTN>. <http://www.dopps.org/annualreport/html/MCOMHTN> (16 July 2016, date last accessed)
62. Agarwal R, Boulidin JM, Light RP *et al.* Inferior vena cava diameter and left atrial diameter measure volume but not dry weight. *Clin J Am Soc Nephrol* 2011; 6: 1066–1072
63. Port FK, Hulbert-Shearon TE, Wolfe RA *et al.* Predialysis blood pressure and mortality risk in a national sample of maintenance hemodialysis patients. *Am J Kidney Dis* 1999; 33: 507–517
64. Salem MM. Hypertension in the haemodialysis population: any relationship to 2-years survival? *Nephrol Dial Transplant* 1999; 14: 125–128
65. Zager PG, Nikolic J, Brown RH *et al.* “U” curve association of blood pressure and mortality in hemodialysis patients. Medical Directors of Dialysis Clinic, Inc. *Kidney Int* 1998; 54: 561–569
66. Hannedouche T, Roth H, Krummel T *et al.* Multiphasic effects of blood pressure on survival in hemodialysis patients. *Kidney Int* 2016; 90: 674–684
67. Mazzuchi N, Carbonell E, Fernandez-Cean J. Importance of blood pressure control in hemodialysis patient survival. *Kidney Int* 2000; 58: 2147–2154
68. Lacson E Jr, Lazarus JM. The association between blood pressure and mortality in ESRD-not different from the general population? *Semin Dial* 2007; 20: 510–517
69. Klassen PS, Lowrie EG, Reddan DN *et al.* Association between pulse pressure and mortality in patients undergoing maintenance hemodialysis. *JAMA* 2002; 287: 1548–1555
70. Foley RN, Herzog CA, Collins AJ. Blood pressure and long-term mortality in United States hemodialysis patients: USRDS Waves 3 and 4 Study. *Kidney Int* 2002; 62: 1784–1790
71. Robinson BM, Tong L, Zhang J *et al.* Blood pressure levels and mortality risk among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. *Kidney Int* 2012; 82: 570–580
72. Cocchi R, Degli EE, Fabbri A *et al.* Prevalence of hypertension in patients on peritoneal dialysis: results of an Italian multicentre study. *Nephrol Dial Transplant* 1999; 14: 1536–1540
73. Atas N, Erten Y, Okyay GU *et al.* Left ventricular hypertrophy and blood pressure control in automated and continuous ambulatory peritoneal dialysis patients. *Ther Apher Dial* 2014; 18: 297–304
74. Cnossen TT, Konings CJ, Fagel WJ *et al.* Fluid state and blood pressure control: no differences between APD and CAPD. *ASAIO J* 2012; 58: 132–136
75. Tonbul Z, Altintepe L, Sozlu C *et al.* Ambulatory blood pressure monitoring in haemodialysis and continuous ambulatory peritoneal dialysis (CAPD) patients. *J Hum Hypertens* 2002; 16: 585–589
76. Enia G, Mallamaci F, Benedetto FA *et al.* Long-term CAPD patients are volume expanded and display more severe left ventricular hypertrophy than haemodialysis patients. *Nephrol Dial Transplant* 2001; 16: 1459–1464
77. Davies SJ. What are the consequences of volume expansion in chronic dialysis patients?: Volume expansion in peritoneal dialysis patients. *Semin Dial* 2015; 28: 239–242
78. Rodby RA, Vonesh EF, Korbet SM. Blood pressures in hemodialysis and peritoneal dialysis using ambulatory blood pressure monitoring. *Am J Kidney Dis* 1994; 23: 401–411
79. Sarafidis PA, Bakris GL. Kidney disease and hypertension. In: G Lip, J Hall, eds. *Comprehensive Hypertension*. London: Mosby Elsevier, 2007, 607–619
80. Hall JE. The renin-angiotensin system: renal actions and blood pressure regulation. *Compr Ther* 1991; 17: 8–17
81. Ritz E, Fliser D. The kidney in congestive heart failure. *Eur Heart J* 1991; 12 (Suppl C): 14–20
82. Machnik A, Neuhofer W, Jantsch J *et al.* Macrophages regulate salt-dependent volume and blood pressure by a vascular endothelial growth factor-C-dependent buffering mechanism. *Nat Med* 2009; 15: 545–552
83. Wiig H, Schroder A, Neuhofer W *et al.* Immune cells control skin lymphatic electrolyte homeostasis and blood pressure. *J Clin Invest* 2013; 123: 2803–2815
84. Dahlmann A, Dorfelt K, Eicher F *et al.* Magnetic resonance-determined sodium removal from tissue stores in hemodialysis patients. *Kidney Int* 2015; 87: 434–441
85. Kelley K, Light RP, Agarwal R. Trended cosinor change model for analyzing hemodynamic rhythm patterns in hemodialysis patients. *Hypertension* 2007; 50: 143–150
86. Karpetas A, Sarafidis PA, Georgianos PI *et al.* Ambulatory recording of wave reflections and arterial stiffness during intra- and interdialytic periods in patients treated with dialysis. *Clin J Am Soc Nephrol* 2015; 10: 630–638
87. Koutroumbas G, Georgianos PI, Sarafidis PA *et al.* Ambulatory aortic blood pressure, wave reflections and pulse wave velocity are elevated during the third in comparison to the second interdialytic day of the long interval in chronic haemodialysis patients. *Nephrol Dial Transplant* 2015; 30: 2046–2053
88. Georgianos PI, Sarafidis PA, Haidich AB *et al.* Diverse effects of interdialytic intervals on central wave augmentation in haemodialysis patients. *Nephrol Dial Transplant* 2013; 28: 2160–2169
89. Georgianos PI, Sarafidis PA, Lasaridis AN. Arterial stiffness: a novel cardiovascular risk factor in kidney disease patients. *Curr Vasc Pharmacol* 2015; 13: 229–238
90. Agarwal R, Light RP. Arterial stiffness and interdialytic weight gain influence ambulatory blood pressure patterns in hemodialysis patients. *Am J Physiol Renal Physiol* 2008; 294: F303–F308
91. Georgianos PI, Agarwal R. Aortic stiffness, ambulatory blood pressure, and predictors of response to antihypertensive therapy in hemodialysis. *Am J Kidney Dis* 2015; 66: 305–312
92. Converse RL Jr, Jacobsen TN, Toto RD *et al.* Sympathetic overactivity in patients with chronic renal failure. *N Engl J Med* 1992; 327: 1912–1918
93. Hausberg M, Kosch M, Harmelink P *et al.* Sympathetic nerve activity in end-stage renal disease. *Circulation* 2002; 106: 1974–1979
94. Ott C, Schmid A, Ditting T *et al.* Renal denervation in a hypertensive patient with end-stage renal disease and small arteries: a direction for future research. *J Clin Hypertens* 2012; 14: 799–801
95. Papademetriou V, Doumas M, Anyfanti P *et al.* Renal nerve ablation for hypertensive patients with chronic kidney disease. *Curr Vasc Pharmacol* 2014; 12: 47–54
96. Desir GV, Wang L, Peixoto AJ. Human renalase: a review of its biology, function, and implications for hypertension. *J Am Soc Hypertens* 2012; 6: 417–426
97. Desir GV. Regulation of blood pressure and cardiovascular function by renalase. *Kidney Int* 2009; 76: 366–370
98. Malyszko J, Koc-Zorawska E, Malyszko JS *et al.* Renalase, stroke, and hypertension in hemodialyzed patients. *Ren Fail* 2012; 34: 727–731
99. Bazzato G, Coli U, Landini S *et al.* Prevention of intra- and postdialytic hypertensive crises by captopril. *Contrib Nephrol* 1984; 41: 292–298
100. Kornerup HJ, Schmitz O, Danielsen H *et al.* Significance of the renin-angiotensin system for blood pressure regulation in end-stage renal disease. *Contrib Nephrol* 1984; 41: 123–127
101. Henrich WL, Katz FH, Molinoff PB *et al.* Competitive effects of hypokalemia and volume depletion on plasma renin activity, aldosterone and catecholamine concentrations in hemodialysis patients. *Kidney Int* 1977; 12: 279–284
102. Lifschitz MD, Kirschenbaum MA, Rosenblatt SG *et al.* Effect of saralasin in hypertensive patients on chronic hemodialysis. *Ann Intern Med* 1978; 88: 23–27
103. Agarwal R, Lewis R, Davis JL *et al.* Lisinopril therapy for hemodialysis hypertension: hemodynamic and endocrine responses. *Am J Kidney Dis* 2001; 38: 1245–1250
104. Drechsler C, Ritz E, Tomaschitz A *et al.* Aldosterone and cortisol affect the risk of sudden cardiac death in haemodialysis patients. *Eur Heart J* 2013; 34: 578–587
105. Abd ElHafeez S, Tripepi G, Mallamaci F *et al.* Aldosterone, mortality, cardiovascular events and reverse epidemiology in end stage renal disease. *Eur J Clin Invest* 2015; 45: 1077–1086

106. Vaziri ND, Ni Z, Wang XQ *et al.* Downregulation of nitric oxide synthase in chronic renal insufficiency: role of excess PTH. *Am J Physiol* 1998; 274: F642–F649
107. Morris ST, McMurray JJ, Spiers A *et al.* Impaired endothelial function in isolated human uremic resistance arteries. *Kidney Int* 2001; 60: 1077–1082
108. Wever R, Boer P, Hijmering M *et al.* Nitric oxide production is reduced in patients with chronic renal failure. *Arterioscler Thromb Vasc Biol* 1999; 19: 1168–1172
109. Lau T, Owen W, Yu YM *et al.* Arginine, citrulline, and nitric oxide metabolism in end-stage renal disease patients. *J Clin Invest* 2000; 105: 1217–1225
110. Schnackenberg CG. Oxygen radicals in cardiovascular-renal disease. *Curr Opin Pharmacol* 2002; 2: 121–125
111. Yokoyama K, Tajima M, Yoshida H *et al.* Plasma pteridine concentrations in patients with chronic renal failure. *Nephrol Dial Transplant* 2002; 17: 1032–1036
112. Mallamaci F, Tripepi G, Maas R *et al.* Analysis of the relationship between norepinephrine and asymmetric dimethyl arginine levels among patients with end-stage renal disease. *J Am Soc Nephrol* 2004; 15: 435–441
113. Vallance P, Leone A, Calver A *et al.* Accumulation of an endogenous inhibitor of nitric oxide synthesis in chronic renal failure. *Lancet* 1992; 339: 572–575
114. Raptis V, Kapoulas S, Grekas D. Role of asymmetrical dimethylarginine in the progression of renal disease. *Nephrology* 2013; 18: 11–21
115. Tada T, Kusano KF, Ogawa A *et al.* The predictors of central and obstructive sleep apnoea in haemodialysis patients. *Nephrol Dial Transplant* 2007; 22: 1190–1197
116. Ogna A, Forni OV, Mihalache A *et al.* Obstructive sleep apnea severity and overnight body fluid shift before and after hemodialysis. *Clin J Am Soc Nephrol* 2015; 10: 1002–1010
117. Abdel-Kader K, Dohar S, Shah N *et al.* Resistant hypertension and obstructive sleep apnea in the setting of kidney disease. *J Hypertens* 2012; 30: 960–966
118. Boyle SM, Berns JS. Erythropoietin and resistant hypertension in CKD. *Semin Nephrol* 2014; 34: 540–549
119. Carlini RG, Dusso AS, Obialo CI *et al.* Recombinant human erythropoietin (rHuEPO) increases endothelin-1 release by endothelial cells. *Kidney Int* 1993; 43: 1010–1014
120. Kang DH, Yoon KI, Han DS. Acute effects of recombinant human erythropoietin on plasma levels of proendothelin-1 and endothelin-1 in haemodialysis patients. *Nephrol Dial Transplant* 1998; 13: 2877–2883
121. Eggena P, Wilsey P, Jamgotchian N *et al.* Influence of recombinant human erythropoietin on blood pressure and tissue renin-angiotensin systems. *Am J Physiol* 1991; 261: E642–E646
122. Hand MF, Haynes WG, Johnstone HA *et al.* Erythropoietin enhances vascular responsiveness to norepinephrine in renal failure. *Kidney Int* 1995; 48: 806–813
123. Koulouridis I, Alfayez M, Trikalinos TA *et al.* Dose of erythropoiesis-stimulating agents and adverse outcomes in CKD: a meta-regression analysis. *Am J Kidney Dis* 2013; 61: 44–56
124. Phrommintikul A, Haas SJ, Elisk M *et al.* Mortality and target haemoglobin concentrations in anaemic patients with chronic kidney disease treated with erythropoietin: a meta-analysis. *Lancet* 2007; 369: 381–388
125. Watson AJ, Gimenez LF, Cotton S *et al.* Treatment of the anemia of chronic renal failure with subcutaneous recombinant human erythropoietin. *Am J Med* 1990; 89: 432–435
126. Krapf R, Hulter HN. Arterial hypertension induced by erythropoietin and erythropoiesis-stimulating agents (ESA). *Clin J Am Soc Nephrol* 2009; 4: 470–480
127. Nissenson AR, Korbet S, Faber M *et al.* Multicenter trial of erythropoietin in patients on peritoneal dialysis. *J Am Soc Nephrol* 1995; 5: 1517–1529
128. Del Vecchio L, Locatelli F. An overview on safety issues related to erythropoiesis-stimulating agents for the treatment of anaemia in patients with chronic kidney disease. *Expert Opin Drug Saf* 2016; 15: 1021–1030
129. Sarafidis PA, Bakris GL. Resistant hypertension: an overview of evaluation and treatment. *J Am Coll Cardiol* 2008; 52: 1749–1757
130. Pullalarevu R, Akbar G, Teehan G. Secondary hypertension, issues in diagnosis and treatment. *Prim Care* 2014; 41: 749–764
131. Kumar A, Hubbard J, Moonim M *et al.* Case report: making the CONN-ec-tion: two cases of persistent hypertension and hypokalaemia following renal transplantation. *Int Urol Nephrol* 2012; 44: 1577–1580
132. Agarwal R, Weir MR. Dry-weight: a concept revisited in an effort to avoid medication-directed approaches for blood pressure control in hemodialysis patients. *Clin J Am Soc Nephrol* 2010; 5: 1255–1260
133. Sinha AD, Agarwal R. Can chronic volume overload be recognized and prevented in hemodialysis patients? The pitfalls of the clinical examination in assessing volume status. *Semin Dial* 2009; 22: 480–482
134. Agarwal R, Andersen MJ, Pratt JH. On the importance of pedal edema in hemodialysis patients. *Clin J Am Soc Nephrol* 2008; 3: 153–158
135. Torino C, Gargani L, Sicari R *et al.* The Agreement between Auscultation and Lung Ultrasound in Hemodialysis Patients: The LUST Study. *Clin J Am Soc Nephrol* 2016; 11: 2005–2011
136. Zhu F, Rosales L, Kotanko P. Techniques for assessing fluids status in patients with kidney disease. *Curr Opin Nephrol Hypertens* 2016; 25: 473–479
137. Zoccali C, Puntorieri E, Mallamaci F. Lung congestion as a hidden threat in end-stage kidney disease: a call to action. *Nephrol Dial Transplant* 2013; 28: 2657–2660
138. Ozkahya M, Toz H, Qzerkan F *et al.* Impact of volume control on left ventricular hypertrophy in dialysis patients. *J Nephrol* 2002; 15: 655–660
139. Kayikcioglu M, Tumuklu M, Ozkahya M *et al.* The benefit of salt restriction in the treatment of end-stage renal disease by haemodialysis. *Nephrol Dial Transplant* 2009; 24: 956–962
140. Cirit M, Akcicek F, Terzioğlu E *et al.* ‘Paradoxical’ rise in blood pressure during ultrafiltration in dialysis patients. *Nephrol Dial Transplant* 1995; 10: 1417–1420
141. Agarwal R, Alborzi P, Satyan S *et al.* Dry-weight reduction in hypertensive hemodialysis patients (DRIP): a randomized, controlled trial. *Hypertension* 2009; 53: 500–507
142. Agarwal R, Bouldin JM, Light RP *et al.* Probing dry-weight improves left ventricular mass index. *Am J Nephrol* 2011; 33: 373–380
143. Zoccali C. Lung water by ultrasound guided treatment in hemodialysis patients (The Lust Study). ClinicalTrials.gov: Bethesda (MD): National Library of Medicine (US). 2016. <https://clinicaltrials.gov/ct2/show/NCT02310061> (2 November 2016, date last accessed)
144. Li PK, Culeton BF, Ariza A *et al.* Randomized, controlled trial of glucose-sparing peritoneal dialysis in diabetic patients. *J Am Soc Nephrol* 2013; 24: 1889–1900
145. Curatola G, Bolignano D, Rastelli S *et al.* Ultrafiltration intensification in hemodialysis patients improves hypertension but increases AV fistula complications and cardiovascular events. *J Nephrol* 2011; 24: 465–473
146. Movilli E, Gaggia P, Zubani R *et al.* Association between high ultrafiltration rates and mortality in uraemic patients on regular haemodialysis. A 5-year prospective observational multicentre study. *Nephrol Dial Transplant* 2007; 22: 3547–3552
147. Weiner DE, Brunelli SM, Hunt A *et al.* Improving clinical outcomes among hemodialysis patients: a proposal for a “volume first” approach from the chief medical officers of US dialysis providers. *Am J Kidney Dis* 2014; 64: 685–695
148. Shimizu K, Kurosawa T, Ishikawa R *et al.* Vasopressin secretion by hypertonic saline infusion during hemodialysis: effect of cardiopulmonary recirculation. *Nephrol Dial Transplant* 2012; 27: 796–803
149. Mc Causland FR, Waikar SS, Brunelli SM. Increased dietary sodium is independently associated with greater mortality among prevalent hemodialysis patients. *Kidney Int* 2012; 82: 204–211
150. Kooman JP, van der Sande F, Leunissen K *et al.* Sodium balance in hemodialysis therapy. *Semin Dial* 2003; 16: 351–355
151. Maduell F, Navarro V. Dietary salt intake and blood pressure control in haemodialysis patients. *Nephrol Dial Transplant* 2000; 15: 2063
152. Tomson CR. Advising dialysis patients to restrict fluid intake without restricting sodium intake is not based on evidence and is a waste of time. *Nephrol Dial Transplant* 2001; 16: 1538–1542
153. Chobanian AV, Bakris GL, Black HR *et al.* The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003; 289: 2560–2572
154. Cybulsky AV, Matni A, Hollomby DJ. Effects of high sodium dialysate during maintenance hemodialysis. *Nephron* 1985; 41: 57–61

155. Ogden DA. A double blind crossover comparison of high and low sodium dialysis. *Proc Clin Dial Transplant Forum* 1978; 8: 157–165
156. Santos SF, Peixoto AJ. Revisiting the dialysate sodium prescription as a tool for better blood pressure and interdialytic weight gain management in hemodialysis patients. *Clin J Am Soc Nephrol* 2008; 3: 522–530
157. Munoz Mendoza J, Sun S, Chertow GM *et al*. Dialysate sodium and sodium gradient in maintenance hemodialysis: a neglected sodium restriction approach? *Nephrol Dial Transplant* 2011; 26: 1281–1287
158. de Paula FM, Peixoto AJ, Pinto LV *et al*. Clinical consequences of an individualized dialysate sodium prescription in hemodialysis patients. *Kidney Int* 2004; 66: 1232–1238
159. Munoz MJ, Bayes LY, Sun S *et al*. Effect of lowering dialysate sodium concentration on interdialytic weight gain and blood pressure in patients undergoing thrice-weekly in-center nocturnal hemodialysis: a quality improvement study. *Am J Kidney Dis* 2011; 58: 956–963
160. Davies S, Carlsson O, Simonsen O *et al*. The effects of low-sodium peritoneal dialysis fluids on blood pressure, thirst and volume status. *Nephrol Dial Transplant* 2009; 24: 1609–1617
161. Tattersall J, Martin-Malo A, Pedrini L *et al*. EBP guideline on dialysis strategies. *Nephrol Dial Transplant* 2007; 22 (Suppl 2): ii5–ii21
162. Kalantar-Zadeh K, Unruh M, Zager PG *et al*. Twice-weekly and incremental hemodialysis treatment for initiation of kidney replacement therapy. *Am J Kidney Dis* 2014; 64: 181–186
163. Kalantar-Zadeh K, Casino FG. Let us give twice-weekly hemodialysis a chance: revisiting the taboo. *Nephrol Dial Transplant* 2014; 29: 1618–1620
164. Obi Y, Streja E, Rhee CM *et al*. Incremental hemodialysis, residual kidney function, and mortality risk in incident dialysis patients: A Cohort Study. *Am J Kidney Dis* 2016; 68: 256–265
165. Ramirez SP, Kapke A, Port FK *et al*. Dialysis dose scaled to body surface area and size-adjusted, sex-specific patient mortality. *Clin J Am Soc Nephrol* 2012; 7: 1977–1987
166. Rayner HC, Zepel L, Fuller DS *et al*. Recovery time, quality of life, and mortality in hemodialysis patients: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis* 2014; 64: 86–94
167. Foley RN, Gilbertson DT, Murray T *et al*. Long interdialytic interval and mortality among patients receiving hemodialysis. *N Engl J Med* 2011; 365: 1099–1107
168. Georgianos PI, Sarafidis PA, Sinha AD *et al*. Adverse effects of conventional thrice-weekly hemodialysis: is it time to avoid 3-day interdialytic intervals? *Am J Nephrol* 2015; 41: 400–408
169. Brunet P, Saingra Y, Leonetti F *et al*. Tolerance of haemodialysis: a randomized cross-over trial of 5-h versus 4-h treatment time. *Nephrol Dial Transplant* 1996; 11 (Suppl 8): 46–51
170. Chertow GM, Levin NW, Beck GJ *et al*. In-center hemodialysis six times per week versus three times per week. *N Engl J Med* 2010; 363: 2287–2300
171. Ok E, Duman S, Asci G *et al*. Comparison of 4- and 8-h dialysis sessions in thrice-weekly in-centre haemodialysis: a prospective, case-controlled study. *Nephrol Dial Transplant* 2011; 26: 1287–1296
172. Kotanko P, Garg AX, Depner T *et al*. Effects of frequent hemodialysis on blood pressure: Results from the randomized frequent hemodialysis network trials. *Hemodial Int* 2015; 19: 386–401
173. Chertow GM, Levin NW, Beck GJ *et al*. Long-term effects of frequent in-center hemodialysis. *J Am Soc Nephrol* 2016; 27: 1830–1836
174. Rocco MV, Daugirdas JT, Greene T *et al*. Long-term effects of frequent nocturnal hemodialysis on mortality: The Frequent Hemodialysis Network (FHN) Nocturnal Trial. *Am J Kidney Dis* 2015; 66: 459–468
175. Daugirdas JT, Greene T, Rocco MV *et al*. Effect of frequent hemodialysis on residual kidney function. *Kidney Int* 2013; 83: 949–958
176. Cice G, Ferrara L, D'Andrea A *et al*. Carvedilol increases two-year survival in dialysis patients with dilated cardiomyopathy: a prospective, placebo-controlled trial. *J Am Coll Cardiol* 2003; 41: 1438–1444
177. Agarwal R, Sinha AD, Pappas MK *et al*. Hypertension in hemodialysis patients treated with atenolol or lisinopril: a randomized controlled trial. *Nephrol Dial Transplant* 2014; 29: 672–681
178. Zannad F, Kessler M, Leheret P *et al*. Prevention of cardiovascular events in end-stage renal disease: results of a randomized trial of fosinopril and implications for future studies. *Kidney Int* 2006; 70: 1318–1324
179. Takahashi A, Takase H, Toriyama T *et al*. Candesartan, an angiotensin II type-1 receptor blocker, reduces cardiovascular events in patients on chronic haemodialysis—a randomized study. *Nephrol Dial Transplant* 2006; 21: 2507–2512
180. Suzuki H, Kanno Y, Sugahara S *et al*. Effect of angiotensin receptor blockers on cardiovascular events in patients undergoing hemodialysis: an open-label randomized controlled trial. *Am J Kidney Dis* 2008; 52: 501–506
181. Iseki K, Arima H, Kohagura K *et al*. Effects of angiotensin receptor blockade (ARB) on mortality and cardiovascular outcomes in patients with long-term haemodialysis: a randomized controlled trial. *Nephrol Dial Transplant* 2013; 28: 1579–1589
182. Tepel M, Hopfenmueller W, Scholze A *et al*. Effect of amlodipine on cardiovascular events in hypertensive haemodialysis patients. *Nephrol Dial Transplant* 2008; 23: 3605–3612
183. Matsumoto Y, Mori Y, Kageyama S *et al*. Spironolactone reduces cardiovascular and cerebrovascular morbidity and mortality in hemodialysis patients. *J Am Coll Cardiol* 2014; 63: 528–536
184. Lin C, Zhang Q, Zhang H *et al*. Long-term effects of low-dose spironolactone on chronic dialysis patients: a randomized placebo-controlled study. *J Clin Hypertens* 2015; 18: 121–128
185. Agarwal R, Sinha AD. Cardiovascular protection with antihypertensive drugs in dialysis patients: systematic review and meta-analysis. *Hypertension* 2009; 53: 860–866
186. Heerspink HJ, Ninomiya T, Zoungas S *et al*. Effect of lowering blood pressure on cardiovascular events and mortality in patients on dialysis: a systematic review and meta-analysis of randomised controlled trials. *Lancet* 2009; 373: 1009–1015
187. Denker MG, Cohen DL. Antihypertensive medications in end-stage renal disease. *Semin Dial* 2015; 28: 330–336
188. Inrig JK. Antihypertensive agents in hemodialysis patients: a current perspective. *Semin Dial* 2010; 23: 290–297
189. Hayashi SY, Seiberger A, Lind B *et al*. Acute effects of low and high intravenous doses of furosemide on myocardial function in anuric haemodialysis patients: a tissue Doppler study. *Nephrol Dial Transplant* 2008; 23: 1355–1361
190. Lemes HP, Araujo S, Nascimento D *et al*. Use of small doses of furosemide in chronic kidney disease patients with residual renal function undergoing hemodialysis. *Clin Exp Nephrol* 2011; 15: 554–559
191. van Olden RW, van Meyel JJ, Gerlag PG. Sensitivity of residual nephrons to high dose furosemide described by diuretic efficiency. *Eur J Clin Pharmacol* 1995; 47: 483–488
192. Medcalf JF, Harris KP, Walls J. Role of diuretics in the preservation of residual renal function in patients on continuous ambulatory peritoneal dialysis. *Kidney Int* 2001; 59: 1128–1133
193. van Olden RW, Guchelaar HJ, Struijk DG *et al*. Acute effects of high-dose furosemide on residual renal function in CAPD patients. *Perit Dial Int* 2003; 23: 339–347
194. Zoccali C, Mallamaci F, Parlongo S *et al*. Plasma norepinephrine predicts survival and incident cardiovascular events in patients with end-stage renal disease. *Circulation* 2002; 105: 1354–1359
195. Jadoul M, Thumma J, Fuller DS *et al*. Modifiable practices associated with sudden death among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. *Clin J Am Soc Nephrol* 2012; 7: 765–774
196. Roberts MA, Pilmore HL, Ierino FL *et al*. The beta-Blocker to Lower Cardiovascular Dialysis Events (BLOCADE) Feasibility Study: A Randomized Controlled Trial. *Am J Kidney Dis* 2016; 67: 902–911
197. Inrig JK, Van BP, Kim C *et al*. Probing the mechanisms of intradialytic hypertension: a pilot study targeting endothelial cell dysfunction. *Clin J Am Soc Nephrol* 2012; 7: 1300–1309
198. Weir MA, Dixon SN, Fleet JL *et al*. beta-Blocker dialyzability and mortality in older patients receiving hemodialysis. *J Am Soc Nephrol* 2015; 26: 987–996
199. Zoccali C, Mallamaci F. Pleiotropic effects of angiotensin II blockers in hemodialysis patients: myth or reality? *Kidney Int* 2014; 86: 469–471
200. London GM, Marchais SJ, Guerin AP *et al*. Salt and water retention and calcium blockade in uremia. *Circulation* 1990; 82: 105–113
201. Aslam S, Santha T, Leone A *et al*. Effects of amlodipine and valsartan on oxidative stress and plasma methylarginines in end-stage renal disease patients on hemodialysis. *Kidney Int* 2006; 70: 2109–2115
202. Pitt B, Rossignol P. Mineralocorticoid receptor antagonists in patients with end-stage renal disease on chronic hemodialysis. *J Am Coll Cardiol* 2014; 63: 537–538

203. Kramann R, Floege J, Ketteler M *et al*. Medical options to fight mortality in end-stage renal disease: a review of the literature. *Nephrol Dial Transplant* 2012; 27: 4298–4307
204. Walsh M, Manns B, Garg AX *et al*. The safety of eplerenone in hemodialysis patients: a noninferiority randomized controlled trial. *Clin J Am Soc Nephrol* 2015; 10: 1602–1608
205. Brest UH. Aldosterone Antagonist Chronic HEModialysis Interventional Survival Trial (ALCHEMIST). *ClinicalTrials.gov*: Bethesda (MD): National Library of Medicine (US); 2000. <https://clinicaltrials.gov/show/NCT01848639> (2 November 2016, date last accessed)

Received: 12.11.2016; Editorial decision: 14.11.2016